



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

101525108
03/09/2003
REC'D 07 OCT 2003
WIPO PCT
#2

Bescheinigung · Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02018210.1

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02018210.1
Demande no:

Anmeldetag: . .
Date of filing: 20.08.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

B.R.A.I.N. Biotechnology Research And
Information Network GmbH
Darmstädter Strasse 34
64673 Zwingenberg
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Isolation and cloning of DNA from uncultivated organisms

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

ISOLATION AND CLONING OF DNA FROM UNCULTIVATED ORGANISMS

EPO - Munich
69
20. Aug. 2002

5 The present invention relates to a device for the isolation and/or purification of
nucleic acid molecules suitable to bind and/or inactivate inhibitors of the activity of
reagents or enzymes used for DNA manipulation and to separate a plurality of
nucleic acid molecules with respect to their size. Moreover, the invention relates to a
method for the isolation of a nucleic acid molecule comprising applying a sample to
10 the device of the invention wherein said nucleic acid molecule preferably is part of a
sample which represents a fraction of the metagenome of a given habitat.
Furthermore, the invention relates to a method for the generation of at least one
gene library comprising nucleic acid molecules isolated by the method of the
invention and to a nucleic acid molecule isolated by the method of the invention and
15 with the device of the present invention.

Several documents are cited throughout the text of this specification. The disclosure
content of the documents cited herein (including any manufacture's specifications,
instructions, etc.) is herewith incorporated by reference.

20

Enzymes are highly efficient biological catalysts. As such they are key players in
environmentally friendly technical conversion processes of modern sustainable
biotechnology.

25

Enzymes particularly from microbial sources are active ingredients in many
processes of the textile, detergent, pulp- and paper, food and feed industries. In
addition widespread stereoselective substrate recognition and conversion make
enzymes particularly attractive for synthetic organic chemists in need of chiral
specificity. A bottleneck in the development of innovative technical processes based
on enzymes is the supply with suitable new biocatalysts. Owing to their phylogeny
and physiological diversity microorganisms constitute the largest resource of natural
30 genetic and enzymatic diversity. However the largest proportion of microorganisms

evades cultivation under laboratory conditions (Amann et. al. (1995). *Microbiol Rev* 59, 143-69). Classic microbiology relying on cultivation of pure strains to provide homogenous and defined systems for homologous enzyme production and to supply genomic DNA for recombinant expression strategies therefore inevitably fails to

5 access the entire biosynthetic potential harboured in this enormous natural resource. The recent development of strategies to directly isolate and clone genomic DNA from non-cultivated microbial consortia opens up new dimensions of accessible enzymatic diversity (Rondon et. al. (2000). *Appl Environ Microbiol* 66, 2541-7). Fundamental work on the handling of DNA from non-cultivated microorganisms - the

10 so-called metagenome – (Handelsman et. al. (1998) *Chem Biol* 5, R245-9) by Torsvik (Torsvik and Goksoyr (1978) *Soil Biology and Biochemistry* 10, 7-12), (Torsvik (1980) *Soil Biology and Biochemistry* 12, 15-21), Somerville (Somerville et. al. (1989) *Appl Environ Microbiol* 55(3), 548-554) and Schmidt (Schmidt et. al. (1991) *Journal of Bacteriology* 173, 4371-4378) showed that genomic DNA can be directly

15 isolated from complex microbial assortments as present, inter alia, in plancton or soil. This DNA may be digested and cloned into suitable vectors for recombinant maintenance in heterologous hosts to generate screenable gene libraries. Such metagenome libraries were shown to be useful in the identification of novel genes from uncultivated organisms. The discovery of novel enzymes by screening of non-

20 normalised metagenome libraries from planctonic and soil sources has been reported in the literature (Cottrell et. al. (1999) *Appl Environ Microbiol* 65, 2553-7), (Henne et. al. (1999) *Appl Environ Microbiol* 65, 3901-7; Henne et. al. (2000) *Appl Environ Microbiol* 66, 3113-3116); (US-patent No : 5,849,491); (Rondon et. al. (2000) *Appl Environ Microbiol* 66, 2541-7). The list of enzyme activities discovered in this

25 way (lipase, esterase, amylase, nuclease, chitinase, xylanase) is still rather small. Importantly also more complex activities like the production of bioactive secondary metabolites requiring entire gene clusters for expression have been identified in metagenomic libraries (MacNeil et. al. (2001) *J Mol Microbiol Biotechnol* 3, 301-8) (Wang et. al. (2000) *Org Lett* 2, 2401-4) (Brady et. al. (2001) *Org Lett* 3, 1981-4).

30 Secondary metabolites, like polyketides, are often produced by enzyme complexes encoded by assortments of genes covering in excess of 100 kbp of contiguous DNA (Schwecke et. al. (1995) *Proc Natl Acad Sci U S A* 92, 7839-43). The cloning of such

large fragments of environmental DNA is much more challenging than cloning smaller DNA fragments and is substantially facilitated by the current invention. Proprietary technology for the cloning particularly of normalised environmental DNA and the screening of libraries generated thereby is described in US patents US 6,280,926; US 6,054,267; US 6,057,103; US 6,001,574 and PCT applications WO99/45154; WO98/58085; WO99/10539.

DNA directly extracted from microbial consortia in the context of their natural substratum usually is contaminated with substances inhibiting standard enzymatic manipulations that are essentially required for cloning, analysis or amplification of nucleic acids carrying genetic information. In particular the efficiencies of DNA digestion with restriction enzymes (Tsai and Olson (1992) *Appl Environ Microbiol* 58, 2292-5), (Tebbe and Vahjen (1993) *Appl Environ Microbiol* 59, 2657-65), the polymerase chain reaction (PCR) (Zhou et. al. (1996) *Appl Environ Microbiol* 62, 316-22), DNA-DNA hybridisation and bacterial transformation with environmental DNA (Tebbe and Vahjen (1993) *loc. cit.*) are inversely correlated with natural substrate derived inhibitor concentrations. Besides inorganic inhibitors like heavy metal ions, there are polysaccharides and in particular humic and fulvic acids that act as the single most important sources of above mentioned inhibitions. Humic and fulvic acids are high molecular weight heterocyclic polyphenols mainly of plant origin with an affinity to polynucleotides and strongly protein denaturing properties ((Young et. al. (1993) *Appl Environ Microbiol* 59, 1972-1974); see appended figure 1).

Yet, the efficient removal of such inhibitors is a prerequisite for all enzymatic manipulations required, e.g., for cloning DNA, in particular environmental DNA, into suitable vectors. Several strategies have been pursued. Simple dilution of contaminated DNA to bring inhibitor concentrations below a critical threshold may be sufficient if the subsequent enzymatic manipulation is of suitable power to compensate for the concomitant reduction in target/substrate concentration. Surely such dilution will significantly curtail the efficiency of most subsequent molecular manipulations necessary for cloning following simple mass- action laws. The polymerase chain reaction (PCR), owing to its exponential amplification strategy is

powerful enough to generate strong signals even from very low target numbers and often reducing the amount of input environmental DNA (and inhibitors) in a reaction will substantially increase the amount of product achieved (Tsai and Olson (1992)

loc. cit.). Gelfiltration of contaminated DNA raw extracts has been used to physically

5 separate DNA from inhibitors based on size differences (Tsai and Olson (1992) loc. cit.), (Jackson et. al. (1997) *Applied and Environmental Microbiology* 63, 4993-4995), (Miller (2001) *J Microbiol Methods* 44, 49-58). Charge differences between DNA and

inhibitors were exploited in strategies using ion-exchange chromatography purification (Tebbe and Vahjen, (1993) loc. cit.); (Straub et. al. (1995) *Water Science*

10 and *Technology* 31, 311-315); (Smalla et. al. (1993) *J Appl Bacteriol*, 74, 78-85). In a

different approach substances showing selective affinity towards polyphenols like soluble polyvinylpyrrolidone (PVP, figure 2, relative molecular weight 10'000-360'000

Da), insoluble polyvinylpolypyrrolidone (PVPP, a crosslinked derivative of PVP) or CTAB (hexadecyltrimethylammonium bromide) have been used to absorb (Holben

15 et. al. (1988) *Appl. Environ. Microbiol.* 54, 703-711) or precipitate inhibitors from solutions (Zhou et. al. (1996) loc. cit.). Berthelet and co-workers used a PVPP

affinity-matrix to chromatograph contaminated DNA solutions on spin columns (Berthelet et. al. (1996) *FEMS Microbiol Lett* 138, 17-22). Using ultracentrifugally

20 generated CsCl density gradients Holben and co-workers (Holben et. al. (1988) loc. cit.) purified DNA from inhibitors based on equilibrium densities. For the construction

of high quality libraries of uniform and particularly large DNA insert sizes (in vectors like Cosmid, Fosmid, BAC) a high resolution size selection step is essential to

provide the reaction with uniformly sized insert DNA, especially if like in the case for BACs the cloning process does not feature any inherent size selective steps. This

25 makes gel electrophoresis particularly attractive for the purification of environmental DNA. Hereby charge-mass ratios and size differences can be exploited

simultaneously to achieve kinetic resolution of DNA from inhibitors and simultaneously the DNA itself is spread out according to size. Although simple gel

electrophoresis may suffice to produce clonable DNA from soils containing only

30 small amounts of humic and fulvic acids (Rondon et. al. (2000) loc. cit.), the humic content of soils varies greatly and can reach up to 60-80 % of the total organic matter (Tsai and Rochelle (2001) *Environmental Molecular Microbiology*, Horizon

Scientific Press, page 15-30 (Extraction of nucleic acids from environmental samples)). Mostly therefore additional purification steps are necessary and still failures to produce clonable soil DNA are common (Entcheva et. al. (2001) Appl Environ Microbiol 67, 89-99).

5

A particular modification of this method was devised by Young and co-workers (Young et. al. (1993) loc. cit.). Here, PVP was added to the gel to selectively lower the charge-mass ratio of humic acids so to improve resolution from DNA. This technique combines an affinity-based selective purification step retarding inhibitors in an electric field with a DNA size resolution step that is indispensable in the preparation of insert DNA for efficient large fragment cloning in vectors like BAC.

10

Yet, prior to subsequent enzymatic modifications of environmental DNA as required for cloning (like PCR using e.g. *Taq* or *Pfu* DNA polymerases or fill-up reactions using e.g. *Klenow*- or T4-DNA-polymerase or ligations using e.g. T4-DNA-ligase or multi-step reactions like phage-packaging) absorbants like PVP must be removed as they themselves are inhibitory for further enzymatic processes.

15

The separation of agarose gel-purified DNA from the PVP absorbant is achieved in the prior art by employing affinity chromatography after melting the DNA containing agarose slice (GeneClean® in (Young et. al. (1993) loc. cit.)). Such a procedure, however, is not suitable to purify very large DNA molecules because shearing forces generated during the elution process will cause fragmentation. Additionally elution efficiency is inversely correlated with molecule size, so large molecules will be selectively lost.

20

Alternatively, large DNA molecules can be electroeluted from an agarose slice cut from a gel after electrophoresis. The DNA will be recovered in diluted form in a buffer like TAE and has to be concentrated before further manipulation. This routinely involves precipitation in 70% ethanol (Ausubel et. al. (Eds.) (1998) Current Protocols in Molecular Biology, John Wiley & Sons, 2.11-2.1.10). Yet, such an alcohol precipitation involves at least one further centrifugation step and, accordingly, adverse shearing forces. Consequently, in order to purify large DNA fragments from agarose gels for enzymatic manipulation and cloning, prior art procedures involve melting a gel slice containing DNA, performing in-gel enzymatic manipulations in a

25

30

re-solidified gel (like end polishing, ligation), solubilizing the gel using a degrading enzyme (Gelase® Epicentre, USA) and transforming the DNA into hosts (*E. coli*).

These procedures are complex and may lead to fragmentation or even loss of nucleic acid molecules. Furthermore, these manipulations of the prior art can not be

5 carried out in the presence of enzyme inhibiting substances like PVP.

Many of the above strategies in different combinations have been used as part of multi-step purification protocols to produce clonable "metagenome DNA". Yet, environmental DNA purification is not trivial and whereas purification of DNA for PCR purposes may be accomplished using commercial kits (FastPrep® *Bio101*, USA), the

10 preparation of sufficient amounts of concentrated and inhibitor-free high-molecular weight DNA (20-300 kbp) for cloning into Cosmids or BACs is much more challenging and may be doomed with failure (Entcheva et. al. (2001) loc. cit).

Thus, the technical problem underlying the present invention was to provide means
15 and method for cloning of genetic material isolated from primary samples. The solution to this technical problem is achieved by providing the embodiments characterized in the claims.

The current invention provides means to overcome the technical difficulties
20 associated with the isolation and cloning of large fragment DNA from uncultivated environmental sources.

Accordingly, the present invention relates to a device for the isolation and/or
purification of nucleic acid molecules comprising at least two layers, a first layer
25 being adapted to bind and/or inactivate inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation and a second layer being adapted to separate a plurality of nucleic acid molecules with respect to their size.

The term "device" as employed herein is an arrangement/construction comprising,
inter alia, gels, or gel chambers or columns as defined herein. Preferably, the gels,
30 gel chambers or columns form the device of the present invention. The nucleic acid molecules to be isolated and/or purified are isolated and/or purified by passing them through the at least two layers of the device as defined herein.

The term "inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation" describes substances comprised in samples of soil, aquatic samples or samples of symbiotic/parasitic consortia which inhibit the activity of reagents or enzymes used in nucleic acid manipulation. Examples of said substances are described herein above and comprise inorganic inhibitors, like, e.g. heavy metal ions, organic inhibitors, like polysaccharides and in particular humic and fulvic acids. Humic and fulvic acids are high molecular weight heterocyclic polyphenols mainly of plant origin with an affinity to polynucleotides and strongly protein denaturing properties. The chemical structure of humic and fulvic acids is shown in appended figures 1A and 1B. Moreover, the chemical properties of said groups of molecules is described in detail by Stevenson (Humus chemistry: genesis, composition, reactions (1994) Wiley New York) and Buffle (Les substances humiques et leurs interactions avec les ions minéraux (1977) Conference Proceedings de la Commission d'Hydrologie Appliquée de A.G.H.T.M.. l'Université d'Orsay, 3-10).

The term "reagents used in nucleic acid manipulation" as used in this context comprises substances like metal ions (e.g. Mg^{2+} , Mn^{2+} , Ca^{2+}), (charged) inorganic and organic molecules required for enzymatic activity or for enzymatic co-factors, said co-factors themselves, or stabilizers. The term "enzymes used in nucleic acid manipulations" relates to enzymes like RNase(s), DNase(s), DNA-polymerase(s), ligase(s) or kinase(s) which are used for nucleic acid manipulation.

The term "nucleic acid manipulation" as used herein comprises standard methods known by the person skilled in the art. Said methods comprise DNA-engineering, such as cloning methods of nucleic acid molecules, the mutation of nucleotide sequences of nucleic acid molecules or amplification methods, like, e.g. PCR. Examples for said methods are described in the appended examples and in laboratory manuals, e.g. Sambrook et. al. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Ausubel et. al. (1998), loc. cit.. In particular, the term "nucleic acid manipulation" relates to the manipulation of DNA or RNA and corresponding cloning techniques.

The term "layer" defines in accordance with the present invention a physical matrix which is characterized by its ability to separate samples containing nucleic acid molecules and also characterized by its ability to separate, if desired, different nucleic acid molecules by their physiological properties, like size or overall charge.

The recited first layer is adapted to characterized by its ability to bind and/or inactivate inhibitors described herein above. The ability of the first layer to bind and/or inactivate inhibitors may be achieved by the addition of compounds with sufficiently high binding affinity to the above described inhibitors so to retard their mobility in aqueous solutions and reduce their effective free concentration so to relieve nucleic acids migrating through the device of the invention from comigrating inhibitors.

The recited second layer is characterized by its ability to separate a plurality of nucleic acid molecules with respect to their size. Accordingly, said physical matrix may be a form of a physical matrix, suited to separate nucleic acid molecules based on molecular sieving, e.g. comprising gels or polymers.

According to a preferred embodiment of the device of invention said first layer is arranged above the second layer.

The term "above" defines the position of the first layer relative to the second layer and relative to the direction in which the samples migrate through the layers of the device. Accordingly, the nucleic acid molecule to be isolated and/or purified with the device of the invention is first contacted with the physical matrix of the first layer and afterwards contacted with the matrix of the second layer. Thus, the present invention comprises devices in which the first layer is horizontally above the second layer as well as devices in which the first layer is vertically above the second layer. Such a device is illustratively exemplified in the appended examples as a gel comprising two phases/layers. Accordingly, as illustrated in the appended figures and examples and described herein, this device may be arranged in form of a gel. Therefore, the device of the invention may be a device, wherein said first layer is a first phase of a gel and said second layer is a second phase of said gel. In its broadest sense, the term "phase" of a gel indicates that this phase has a different overall chemical constitution

than a further (second) phase of said gel. For example, the difference in the overall chemical construction may be due to the presence of chemicals/compounds that bind or inactivate the above mentioned inhibitors of the activity of reagents or enzymes used in nucleic acid manipulations.

5 Preferably, the device of the present invention comprises a gel, wherein said gel is an agarose-gel or a polyacrylamid-gel.

Methods for the preparation of said gels are known by the person skilled in the art and are described in the appended examples and in standard laboratory manuals, e.g. Sambrook et. al. (1989), loc. cit., Cold Spring Harbor, New York; Ausubel et. al. (1998), loc. cit..

10 Preferably, the device of the invention comprises in said first layer polyvinylpyrrolidone (PVP), or polyvinylpolypyrrolidone (PVPP) or combinations thereof. As demonstrated in the appended examples PVP and PVPP are immobile components of said first layer or are characterized at least by a lower mobility compared to the nucleic acid molecules and to bind or interact with the above characterized "inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation".

20 Further examples of corresponding components of said first layer (i.e. inactivators of the inhibitors defined herein above) are functional molecules like CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, nucleic acids immobilized or tethered on appropriate matrices acting as catcher molecules or ion-exchanger. These functional molecules may act by complexing ions (like EDTA, EGTA,), precipitating polysaccharides (like CTAB), binding small hydrophobic molecules or uncharged small molecules, like cyclodextrins, binding specific surface structures (like proteins, (poly)peptides and aptamers). Said proteins may comprise, e.g., antibodies and (poly)peptides directed against inhibitors. Furthermore, lectins are envisaged as inactivators of the inhibitors defined herein. An example for a nucleic acids acting as catcher molecules in accordance with the invention are RNA-aptamers.

30 The term "(poly)peptide" as used herein summarizes a group of molecules which comprise the group of peptides, consisting of up to 30 amino acids, as well as the

group of polypeptides, consisting of more than 30 amino acids.

Above described low molecular weight compounds comprise electrically charged compounds (e.g. CTAB, EDTA) and uncharged small molecules (e.g. cyclodextrin).

Said compounds are soluble in aqueous solutions and, thus, mobile due to rapid
5 diffusion and in particular mobile in an electrical field. In order to achieve a lower mobility of said compounds compared to the nucleic acid molecules, said compounds may be coupled to the physical matrices (e.g. chemically coupled). Compounds which are soluble in aqueous solutions but not electrically charged (e.g. PVP) and compounds which are non-soluble in aqueous solutions (e.g. PVPP) show
10 a lower mobility compared to the nucleic acid molecules and thus do not essentially require to be coupled to the physical matrices.

More preferably, the device the invention is a device, wherein said second layer is substantially free of PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins,
15 (poly)peptides, nucleic acids or ion-exchanger.

The term "substantially free" is understood in accordance with the invention to define a layer which does not contain the recited compounds in an effective amount which is detectable by standard methods. Said standard methods are known in the art and comprise MS (mass spectrometry), FT-IR (fourier transform infrared spectrometry),
20 NMR (nuclear magnetic resonance) or HPLC (high performance liquid chromatography). Accordingly, the second layer does, most preferably, not contain any of the recited compounds as an essential/effective element.

According to a more preferred embodiment the device of the invention is electrically
25 biased to enhance flow of (a) sample(s) through the layers.

As known by the person skilled in the art nucleic acid molecules are negatively charged due to the ribose-phosphate framework. Accordingly, nucleic acid molecules migrate in an electrical field from the cathode (-) to the anode (+).

Examples for devices which are electrically biased to enhance flow of (a) sample(s)
30 are devices for gel electrophoresis. Devices with continuous as well as devices with discontinuous electrical fields are particularly comprised by the present invention. Accordingly, the electrical field can be discontinuous due to a gradient of a salt

(buffer salt), a pulsed field comprising e.g. different angles. Again, an electrically biased device in accordance with the present invention is shown in the appended figures and illustrated in the appended examples.

- 5 The invention relates to a device, wherein said first layer preferably comprises sample loading means.

10 The term "sample loading means" defines in accordance with the invention means for placing the sample comprising nucleic acid molecules in the device of the invention. Examples for said means are sample slots in a gel, the surface of the matrix of a column or a valve for injecting a sample onto a column.

The isolation of nucleic acid molecules by using a device of the invention in the format of a horizontal agarose-gel is described in the appended example 1 and in the format of a single column is described in the appended example 4. Examples for the corresponding devices are shown in the appended figures 3 and 4.

15 In a more preferred embodiment of the invention said loading means are provided in an array in an upper portion of the first layer, defining an array of columns, each being capable of isolating nucleic acid molecules.

20 Hence, a device according to the preferred embodiment of the invention comprises more than one means for placing the sample. Thus, it is possible to isolate nucleic acid molecules from different samples in parallel. An example for loading means provided in an array is a gel comprising different loading slots and, therefore, different lanes (lines in which the samples are separated). An alternative example for said array are columns which are arranged in groups, e.g. in a frame. Said frame
25 may comprise low numbers of columns (two to twelve) for the isolation of nucleic acid molecules from low numbers of samples. Also in accordance with the invention are frames comprising medium numbers of columns (twelve to 96) as well as frames with high numbers of columns (more than 96) which are suitable for high throughput screens (HTS).

30 According to an alternative embodiment of the invention said first layer of the device is arranged below the second layer.

The term "below" defines the position of the second layer relative to the first layer and relative to the direction in which the samples migrate through the layers of the device. The present invention comprises devices for the isolation and/or purification

of nucleic acid molecules in which the nucleic acid molecules are first contacted with the physical matrix of the second layer to separate the molecules with respect to their size and subsequently contacted with the physical matrix of the first layer to bind and/or inactivate the above defined inhibitors.

Accordingly, the present invention comprises devices in which the second layer is horizontally above the first layer as well as devices in which the second layer is vertically above the first layer. Most preferred is in this context a device in form of a column, wherein said column comprises said first and said second layer.

An alternatively preferred embodiment of the invention relates to a device, wherein said second layer is a first phase of a column and said first layer is a second phase of said column. As pointed out herein above, the first layer comprises functional molecules capable of inactivating or binding inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation.

It is also envisaged, in accordance with the invention, that an enhancement of the flow of (a) sample(s) through the layers of the device is, for example, achieved by gravity or by the pressure of the flow of a diluent. Examples for appropriate diluents are buffer solutions. The appended Examples show a device in form of a column as described herein.

Preferably, the device of the invention comprises in said first layer (in a column preferably the second phase) a matrix comprising PVP or PVPP or combinations thereof. As described herein above PVP and PVPP are characterized at least by a lower mobility compared to the nucleic acid molecules and to bind or interact with the above characterized "inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation".

Further examples of corresponding components of said first layer (i.e. compounds capable of inactivating the inhibitors of nucleic acid manipulations) are EDTA, EGTA, CTAB, cyclodextrins, proteins, (poly)peptides, nucleic acids acting as catcher

molecules or ion-exchangers. Said proteins may comprise, e.g., antibodies directed against inhibitors. Also lectins are envisaged as inactivators. An example for said nucleic acids acting as catcher molecules in accordance with the invention are RNA-aptamers.

5

More preferably the device the invention is a device, wherein said second layer (in a column preferably the first phase) is a matrix which is substantially free of PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, nucleic acids or ion-exchangers.

10

The term "substantially free" is defined herein above.

According to a further preferred embodiment of the invention, said matrix of said first and/or second layer is agarose, sepharose™, sephadex™, sephacryl™, BioGel™, superose™ or acrylamide.

15

Variations of the samples comprising the nucleic acid molecules to be isolated and/or purified with the device of the invention may require specific materials of the matrix of the first and/or the second layer. Said variations may concern the quality of the comprised nucleic acid molecules as well as the quality of the sample itself with respect to characteristic substances which may be contained. Accordingly, the matrix of the first and/or the second layer may be a specific matrix for gel-filtration which allows a molecular sieving of the nucleic acid molecules. Suitable media for matrices are characterized by a specific size of the pores which is known by the person skilled in the art and described in the instructions provided by the manufacturer of commercially available media. Said matrices comprise media for gel-filtration. Examples of said media comprise sepharose™ (e.g. sepharose2B, sepharose4B, sepharose6B), sephadex™ (e.g. sephadex G200, sephadex G150) sephacryl™ and superose™ as well as bio-gel™ P100 and bio-gel™ P200. Moreover, said media comprise agarose, polymers as e.g. dextrans and acylamid based-resins. In particular, media are preferred which are suitable for two-phases-columns. Further suitable materials are known to the person skilled in the art.

20

25

30

In a further preferred embodiment of the device of the invention said nucleic acid molecule is DNA or RNA. Most preferably, said DNA is genomic DNA (gDNA).

A particularly preferred embodiment of the invention is a device, wherein said nucleic acid molecule is derived from (micro)organisms of soil, sediments, water, for example sea water, or symbiotic/parasitic consortia.

The term "soil" defines in accordance with the invention the complex product of geological and biological processes acting on inorganic minerals and biomass deposited on the earth surface. It contains the majority of microbial biodiversity on earth. (Whitman et. al. (1998) Proc Natl Acad Sci USA, 95(12, 6578-83) acting to recycle and biomineralize organic matter and serves as a substratum to anchor and nourish higher plants.

In the appended examples the preparation of nucleic acid molecules isolated from soil is exemplarily described.

Examples for symbiotic/parasitic consortia in accordance with the present invention are consortia isolated e.g. from animal tissues or organs, e.g. from gut, stomach, intestines, like appendix or insect-, bird- and mammalian-intestinal tracts or -guts, comprising ruminant-gut. Also envisaged are animal tissues or organs from annelid(s), mollusc(s), sponge(s), cnidaria, arthropod(s), amphibian(s), fish or reptile(s). However, it is also envisaged that nucleic acids from parasitic consortia from human tissue, organs, sputum, faeces, sperm, blood, urine or other body fluids are isolated.

More preferably said (micro)organisms from which said nucleic acid molecules are derived from are (micro)organisms of aquatic plancton, animal tissues and organs as described herein above, microbial mats, clusters, sludge flocs, or biofilms.

(Micro)organisms of the "aquatic plancton" comprises bacterial plancton, archaeal plancton, viruses, phytoplankton as well as zooplankton. Said (micro)organisms are known as small organisms.

Biofilms are microbial assemblages on a surface in "aqueous environments" in which microbes are embedded in a hydrated polymeric matrix. This matrix acts like a kind of glue, holding the microbes together, attaching them to the surface and protecting

them from detrimental external influences. They may contain several taxonomically distinct species (e.g. bacteria, fungi, algae, and protozoa) and may form on surfaces of diverse composition like e.g. metals, glass, plastics, tissue, mineral and soil particles.

- 5 Microbial mats and clusters are microbial assemblages/aggregates similar to biofilms in composition however not necessarily as firmly attached to solid surfaces.

10 According to a further preferred embodiment of the invention said (micro)organisms are isolated and/or purified as consortia of coexisting species. Preferably it is envisaged that said (micro)organisms are isolated and/or purified as consortia of coexisting species without previous separation/purification of single microorganismic species.

15 Preferably said consortia of coexisting species comprise at least one organism that cannot proliferate indefinitely in an artificial setting (e.g. a synthetic or semisynthetic culture medium) in the absence of other species and/or in the absence of the substratum it is isolated from, and wherein said substratum contains the above defined inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation.

20 To obtain the DNA of said microorganisms they may either be bulk-separated mechano-chemically from most of the surrounding matrix they are attached to or embedded or suspended in (like water, soil, sediments, organic debris of plant or animal origin) before lysis (indirect lysis) or they may be lysed and extracted directly i.e. still in the context of/attached to their surrounding physical matrix (soil, biofilm, floc, cluster) that contains a plurality of potential inhibitors of molecular, in particular enzymatic manipulation. Bulk-separation of microorganisms may be accomplished e.g. through mechanical agitation, ion-exchange resin mediated desorption (optionally facilitated through substances added to the extraction buffers used like detergents, salts) followed by an optional concentration step like filtration (for suspended plankton) or suspending and differential gravitational sedimentation in a liquid. In both instances total nucleic acids of mixed origin are isolated irrespective of taxonomic status, abundance and cultivability of the respective taxonomically mixed species.

Classic isolation of nucleic acid molecules from single microbial species or groups of microbial species comprise cultivation of said organisms by massive dilution in selectively growth supporting media. Thereby above mentioned inhibitors of

molecular manipulation of nucleic acids are diluted to facilitate subsequent isolation and cloning. Yet at the cost of a massive reduction in species representation as very few species can be supported by standard cultivation techniques (see herein above). In contrast the present invention provides means for the isolation of nucleic acid molecules derived from taxonomically mixed (micro)organisms without cultivation in synthetic media. Since such steps of cultivation under laboratory conditions results in depletion or at least in significant dilution of organisms which do not grow under said conditions the present invention surprisingly provides means for the isolation and/or purification of nucleic acid molecules derived from said organisms.

In a preferred embodiment of the invention said nucleic acid molecules represent a fraction of the metagenome of a given habitat.

As known in the art the term "metagenome" defines the totality of all genomes of organisms of a given habitat and is furthermore defined in the art; see inter alia Handelsman et. al. (1998) loc. cit.. In particular, the term "metagenome" relates to genomic nucleic acids, preferably DNA, derived from unknown or uncultivable microorganisms, i.e. organisms that cannot be isolated by standard methods and made actively replicating in standard artificial media for indefinite periods of time.

Accordingly the term "a fraction of the metagenome of a given habitat" defines in accordance with the invention nucleic acid molecules and in particular large nucleic acid molecules (>200bp) derived from the total pool of heterogenous microbial genomes present in a given habitat. This is irrespective of phylogenetic affiliation or molecular or physiological traits. Particularly the representation of any particular microbial genome in the extracted portion of the metagenome is not influenced by or dependent on the cultivatability of this organism. Therefore nucleic acids of uncultivated and in a preferred form particularly of uncultivable (micro)organisms are substantially represented in the extracted fraction of the metagenome.

An alternative embodiment of the invention relates to a method for the isolation of a nucleic acid molecule comprising applying a sample to the device as defined herein above. Said sample (for example derived from soil, sediments, water or symbiotic/parasitic consortia) is loaded onto a device comprising the above-identified at least two layers (for example a gel or a column comprising said two layers) and the nucleic acid molecule is purified and/or isolated by passing them through said two layers. In accordance with the invention, the first layer as defined herein, i.e. the layer comprising the inactivators of the inhibitors of reagents and enzymes of nucleic acid manipulations, retains said inhibitors, wherein said second layer provides for, e.g., an isolation and separation step for isolating/separating the nucleic acid molecules in accordance with their physical properties, like size or overall charge. It is envisaged and documented in the appended examples, that the nucleic acid molecules to be isolated pass completely through the two layers of the device of the invention (e.g. a column comprising the two layers) or that the nucleic acid molecules pass only partially through the second layer. For example, it is envisaged that said nucleic acid molecules pass completely through said first layer comprising the inactivators of inhibitors and only partially through said second layer which is substantially free of said inactivators. This method is, inter alia, employed in the device of the invention which is in form of a gel. The nucleic acid molecules to be isolated and/or purified may be isolated or purified from the second layer by conventional means, for example by electroelution.

The method of the invention is also exemplified in the appended examples 1, 4 and 5.

Said method may optionally comprise one or more additional steps of subsequent purification of the obtained nucleic acid molecule(s).

According to a preferred embodiment of the method of the invention a fractions of the metagenome is isolated from a given habitat.

The invention relates in an alternative embodiment to a method for the generation of at least one gene library, comprising the steps of

- (a) isolating and/or purifying nucleic acid molecules from a sample using a device as defined herein above and optionally amplifying said nucleic acid molecules;
- (b) cloning the isolated and/or purified and optionally amplified nucleic acid molecules into appropriate vectors; and

- 5 (c) transforming suitable hosts with said suitable vectors.

Methods for the amplification of an isolated and/or purified nucleic acid molecule are known in the art and comprise e.g. polymerase chain reaction (PCR).

10 Methods for the cloning of nucleic acid molecules into appropriate vectors and transforming suitable hosts with said suitable vectors represent standard methods of molecular biology and are described in the appended examples (in particular, example 2) and in laboratory manuals, e.g. Sambrook et. al. (1989) loc. cit.; Ausubel et. al. (1998), loc. cit.; Mülhardt (2002) Der Experimentator: Molekularbiologie/Genomics; Gustav Fischer. Suitable vectors are described herein below.

15 The above described method may, optionally, additionally comprise one or more of the following steps prior to the cloning of nucleic acid molecules into appropriate vectors according to step (b):

- (i) modification of DNA ends of the isolated and/or purified nucleic acid molecules, e.g. to remove or fill-up random 3'- or 5'-overhangs (polishing/fill-up/blunting), by DNA polymerase treatment (e.g. with Klenow enzyme, T4-
20 polymerase) or treatment with exonucleases (like mung bean nuclease) or introducing defined 3'overhangs (like single nucleotide overhangs, in particular adenosin overhangs) using, e.g. DNA polymerase (like Taq), for subsequent cloning into appropriate vectors (e.g. T-overhang vectors like pGEM T-easy from Promega, USA) or introducing 3'- or 5'-overhangs using restriction endo-
25 nucleases;
- (ii) phosphorylation of the isolated and/or purified nucleic acid molecules (e.g. by PNK); and/or
- (iii) ligation of the isolated and/or purified nucleic acid molecules to other nucleic acid molecules by treatment with an enzymatic ligase (e.g. by T4-ligase) or
30 topoisomerase.

Also envisaged is a "sizing" step, as also illustrated briefly in (i) herein above, wherein said step comprises sizing the obtained nucleic acid molecules by treatment

of the isolated and/or purified nucleic acid molecules with an enzymatic endonuclease (e.g. by restriction endonucleases or DNase I) and/or mechanical shearing (e.g. by ultrasonication or passing nucleic acids with high pressure through narrow tubes or valves similar to the "nebulizer" from Invitrogen, USA).

5 Suitable vectors comprise plasmid vectors (e.g. pUC18 and derivatives thereof, pBluescript etc.), cosmid vectors (e.g. Expand, SuperCos), fosmid vectors (e.g. EpiFos 5), phage vectors (e.g. lambda ZAP), BAC vectors (e.g. pBeloBAC) and YAC vectors.

10 Preferably, said suitable hosts are selected from the group consisting of *E. coli*, *Pseudomonas* sp., *Bacillus* sp, *Streptomyces* sp, other actinomycetes, myxobacteria yeasts and filamentous fungi.

Transformation of said suitable host may be assayed by standard methods of molecular biology; see Sambrook et. al. (1989) loc. cit.; Ausubel et. al. (1998), loc. cit.; Mülhardt (2002) loc. cit.. Corresponding assays for a successful transformation may be based on sequence similarity and performed on plated colonies (filter hybridization) by probe hybridization or by PCR analysis of colonies or extracted DNA of single or multiple colonies. Methods for the preparation of suitable oligonucleotides are known in the art. Alternatively the inserts may be sequenced and targets identified via homology search in appropriated sets of deposited sequence data (e.g. GenBank).

20 Activity based assays may be performed by screening for substrate conversion/degradation inside or outside the host (scoreable by substrate clearing zones around colonies; color development; molecular product profiling by high performance liquid chromatography (HPLC), mass spectrography (MS) or gas chromatography (GC); complementation of growth-deficient mutants), by screening for growth inhibition/stimulation of indicator organisms (in overlays).

25 Accordingly, and in a further embodiment, the present invention relates to a gene library obtained by the method disclosed herein and by employing the device described in this invention.

An alternative embodiment of the invention relates to a gene library generated from metagenome nucleic acid molecules, preferably from DNA, from non-planctonic (micro)organisms comprising average insert sizes of at least 50 kB, at least 55 kB, at least 60 kB, at least 70 kB, at least 80 kB, at least 90 kB or at least 100 kB.

5 As defined herein above planctonic (micro)organisms of the "aquatic plancton" comprises bacterial- and archaeal plancton, viruses, phytoplankton as well as zooplankton. Said (micro)organisms are known as small organisms living in aquatic habitats. Accordingly the term "non-planctonic (micro)organisms" defines (micro)organisms in accordance with the invention which are not comprised by the
10 term "planctonic (micro)organisms". This group of (micro)organisms comprise (micro)organisms of soil, microbial mats, clusters sludge flocs, biofilms and symbiotic/parasitic consortia.

Similarly, the invention relates to a gene library generated from metagenome nucleic
15 acid molecules, preferably DNA, from planctonic (micro)organisms comprising average insert sizes of at least 85 kB, at least 90 kB, at least 95 kB, at least 100 kB, at least 120 kB, at least 140 kB, at least 160 kB or at least 200 kB.

In contrast to the group of "non-planctonic (micro)organisms" defined herein above the term "planctonic (micro)organisms" defines (micro)organisms of the "aquatic
20 plancton" comprises bacterial- and archaeal plancton, viruses, phytoplankton and zooplankton as described herein above.

The average insert size of a (gene) library is, inter alia, determined by (a) isolating the cloned recombinant DNA of at least 0.1% of the clones of the respective library
25 (however no less than a minimum of 20 clones) by methods known in the art, (b) digesting the isolated cloned recombinant DNA molecules with restriction enzymes (6-base or 8-base cutters, e.g., BamHI or EcoRI or Not I used singular or in combination) so to preferentially digest the vector backbone away from insert DNA (e.g. Not I used for pEpiFos5), (c) separating the resulting DNA fragments obtained
30 from each clone individually by agarose gel electrophoresis (continuous or pulsed-field) as known in the art and (d) adding together the sizes of all non-vector

fragments of all analyzed clones of a library and dividing the resulting number by the number of clones analyzed in order to obtain a figure for the average insert size.

The present invention relates further to a nucleic acid molecule comprising a DNA as depicted in SEQ ID NO: 1 or comprising a DNA as deposited under EMBL accession number AJ496176.

Said nucleic acid molecule of the invention has been isolated and obtained by employing the device of the invention. It represents a part of the genome of the newly identified Crenarchaeote as isolated with methods described herein and by techniques implied in the device of the present invention. Taxonomically the crenarchaeota represent a prokaryotic phylum as part of the archeal kingdom. The majority of its representatives are known to be hyperthermophiles yet increasingly they are found in mesophilic habitats as well; see Burggraf et al. (1997) Int J Syst Bacteriol., 47, 657-660; Preston et al. (1996) Proc Natl Acad Sci USA., 93, 6241-46.

In the context of the present invention, the term "genome" defines not only sequences which are open reading frames (ORFs) encoding proteins, polypeptides or peptides, but also refers to non-coding sequences. Accordingly, the term "nucleic acid molecule" comprises coding and, wherever applicable, non-coding sequences. The nucleic acid molecule of the invention furthermore comprises nucleic acid sequences which are degenerative to the above nucleic acid sequences. In accordance with the present invention, the term "nucleic acid molecule" comprises also any feasible derivative of a nucleic acid to which a nucleic acid probe may hybridize. Said nucleic acid probe itself may be a derivative of a nucleic acid molecule capable of hybridizing to said nucleic acid molecule or said derivative thereof. The term "nucleic acid molecule" further comprises peptide nucleic acids (PNAs) containing DNA analogs with amide backbone linkages (Nielsen, P., Science 254 (1991), 1497-1500). The term "ORF" ("open reading frame") which encodes a polypeptide, in connection with the present invention, is defined either by (a) the specific nucleotide sequences encoding the polypeptides specified above in (aa) or in (ab) or (b) by nucleic acid sequences hybridizing under stringent conditions to the complementary strand of the nucleotide sequences of (a) and encoding a polypeptide deviating from the polypeptide of (a) by one or more amino acid

substitutions, deletions, duplications, insertions, recombinations, additions or inversions.

Furthermore the present invention relates in one embodiment to a nucleic acid molecule representing part of the genome of a non-thermophilic crenarchaeote, whereby said nucleic acid molecule has at least one of the following features:

- (a) it contains at least one ORF which encodes a polypeptide having the amino acid sequence SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35;
- (b) comprises the DNA sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;
- (c) it comprises portion of at least 20 nucleotides, preferably 100 nucleotides, more preferably at least 500 nucleotides which hybridize under stringent conditions to the complementary strand of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;
- (d) it is degenerate as a result of the genetic code with respect to the nucleic acid molecule of (c); or
- (e) it is at least 50% identical with the nucleic acid molecule of SEQ ID NO: 2, SEQ ID NO: 20 or SEQ ID NO: 30, 45% identical with the nucleic acid molecule of SEQ ID NO: 8 or SEQ ID NO: 26, 35% identical with the nucleic acid molecule of SEQ ID NO: 16, SEQ ID NO: 22 or SEQ ID NO: 24 or 30% identical with the nucleic acid molecule of SEQ ID NO: 4, SEQ ID NO: 14, SEQ ID NO: 18 or SEQ ID NO: 28;

The following list relates to the SEQ ID NOS as defined herein and shows (partial) identification of ORFs as defined herein:

Name	Identity
SEQ ID NO:1	complete DNA sequence 29i4, 33925 nt, of a newly identified Crenarchaeote
SEQ ID NO:2	ORF001 DNA sequence, 2367 nt, Fam. B DNA Polymerase (truncated ORF)
SEQ ID NO:3	ORF001 Protein sequence, 789 aa, Fam. B DNA Polymerase (truncated protein)
SEQ ID NO:4	ORF002 DNA sequence, 882 nt, α/β hydrolase
SEQ ID NO:5	ORF002 Protein sequence, 293 aa, α/β hydrolase
SEQ ID NO:6	ORF003 DNA sequence, 318 nt
SEQ ID NO:7	ORF003 Protein sequence, 105 aa
SEQ ID NO:8	ORF004 DNA sequence, 1086 nt, Polyhydroxyalkanoate Synthase
SEQ ID NO:9	ORF004 Protein sequence, 361 aa, Polyhydroxyalkanoate Synthase
SEQ ID NO:10	ORF005 DNA sequence, 582 nt
SEQ ID NO:11	ORF005 Protein sequence, 193 aa
SEQ ID NO:12	ORF006 DNA sequence, 438 nt
SEQ ID NO:13	ORF006 Protein sequence, 145 aa
SEQ ID NO:14	ORF007 DNA sequence, 915 nt, Glycosyl Transferase group 1
SEQ ID NO:15	ORF007 Protein sequence, 304 aa, Glycosyl Transferase group 1
SEQ ID NO:16	ORF008 DNA sequence, 1692 nt, Asparagine Synthase
SEQ ID NO:17	ORF008 Protein sequence, 563 aa, Asparagine Synthase
SEQ ID NO:18	ORF009 DNA sequence, 666 nt, Phosphoserin Phosphatase
SEQ ID NO:19	ORF009 Protein sequence, 221 aa, Phosphoserin Phosphatase
SEQ ID NO:20	ORF010 DNA sequence, 1212 nt
SEQ ID NO:21	ORF010 Protein sequence, 403 aa
SEQ ID NO:22	ORF011 DNA sequence, 1164 nt, Transmembrane protein
SEQ ID NO:23	ORF011 Protein sequence, 387 aa, Transmembrane protein

SEQ ID NO:24	ORF012 DNA sequence, 882 nt, Fix A Electron Transfer Flavoprotein
SEQ ID NO:25	ORF012 Protein sequence, 293 aa, Fix A Electron Transfer Flavoprotein
SEQ ID NO:26	ORF013 DNA sequence, 1284 nt, Fix B Electron Transfer Flavoprotein
SEQ ID NO:27	ORF013 Protein sequence, 427 aa, Fix B Electron Transfer Flavoprotein
SEQ ID NO:28	ORF014 DNA sequence, 1878 nt, Fix CX Fusion Electron Transfer Flavoprotein
SEQ ID NO:29	ORF014 Protein sequence, 625 aa, Fix CX Fusion Electron Transfer Flavoprotein
SEQ ID NO:30	ORF015 DNA sequence, 2238 nt, Sensory Transduction Histidin Kinase
SEQ ID NO:31	ORF015 Protein sequence, 745 aa, Sensory Transduction Histidin Kinase
SEQ ID NO:32	ORF016 DNA sequence, 519 nt
SEQ ID NO:33	ORF016 Protein sequence, 172 aa
SEQ ID NO:34	ORF017 DNA sequence, 1008 nt, (truncated ORF)
SEQ ID NO:35	ORF017 Protein sequence, 335 aa, (truncated protein)

A potential field of application of "ORF004" as defined herein comprise the generation or modification of biogenic polymers/polyesters (polyhydroxyalkanoate synthase, Zinn, (2001) Adv Drug Deliv Rev 53(1):5-21 and Fidler, (1992), FEMS Microbiol Rev 9(2-4):231-5; Snell, (2002) Metab. Eng. 4(1):29-40)

Furthermore, the ORF008, a potential asparagine synthetase may be used in the context of amino acid synthesis (EC 6.3.5.4) and/or for the generation of transgenic organisms, like bacteria, plants with altered capacities to generate amino acids.

10

In addition, ORFs 12, 13 or 14 may play a role in redox processes involved in nitrogen fixation and may be useful in generating transgenic organisms like bacteria,

plants with altered capacities to assimilate nitrogen.

The term "hybridizing" as used herein refers to a pairing of polynucleotides to a complementary strand of polynucleotide which thereby form a hybrid. Said complementary strand polynucleotides are, e.g. the polynucleotides of the invention or parts thereof. Therefore, said polynucleotides may be useful as probes in Northern or Southern Blot analysis of RNA or DNA preparations, respectively, or can be used as oligonucleotide primers in PCR analysis dependent on their respective size. Preferably, said hybridizing polynucleotides comprise at least 10, more preferably at least 15 nucleotides in length while a hybridizing polynucleotide of the present invention to be used as a probe preferably comprises at least 100, more preferably at least 200, or most preferably at least 500 nucleotides in length.

It is well known in the art how to perform hybridization experiments with nucleic acid molecules, i.e. the person skilled in the art knows what hybridization conditions s/he has to use in accordance with the present invention. Such hybridization conditions are referred to in standard text books such as Sambrook et. al. (1989) loc. cit. or Higgins, S.J., Hames, D. "RNA Processing: A practical approach", Oxford University Press (1994), Vol. 1 and 2.

"Stringent hybridization conditions" (also referred to highly stringent conditions as contrasted to conditions of low stringency) refers to conditions which comprise, e.g. an overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 x SSC at about 65°C. Said conditions for hybridization are also known by a person skilled in the art as "high stringent conditions for hybridization". Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the invention at lower stringency hybridization conditions ("low stringent conditions for hybridization"). Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37°C in a solution comprising 6X SSPE

(20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1 X SSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes

performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC). Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Preferred in accordance with the present inventions are polynucleotides which are capable of hybridizing to the polynucleotides of the invention or parts thereof, under stringent hybridization conditions, i.e. which do not cross hybridize to unrelated polynucleotides.

The nucleic acid molecules that are homologous to the above-described molecules and that represent derivatives of these molecules usually are variations of these molecules that represent modifications having the same biological function. They can be naturally occurring variations, for example sequences from other organisms, or mutations that can either occur naturally or that have been introduced by specific mutagenesis. Furthermore, the variations can be synthetically produced sequences.

The allelic variants can be either naturally occurring variants or synthetically produced variants or variants produced by recombinant DNA processes.

Generally, by means of conventional molecular biological processes it is possible (see, e.g., Sambrook et. al. (1989) loc. cit.) to introduce different mutations into the nucleic acid molecules of the invention. One possibility is the production of deletion mutants in which nucleic acid molecules are produced by continuous deletions from the 5'- or 3'-terminus of the coding DNA sequence and that lead to the synthesis of proteins that are shortened accordingly. Another possibility is the introduction of single-point mutation at positions where a modification of the amino acid sequence influences, e.g., the enzyme activity or the regulation of the enzyme. By this method muteins can be produced, for example, that possess a modified K_m-value or that are

no longer subject to the regulation mechanisms that normally exist in the cell, e.g. with regard to allosteric regulation or covalent modification. Such muteins may be identified, e.g. by methods of the present invention, to be valuable as therapeutically useful modulators (inhibitors/antagonists or enhancer/agonists) of the activity of the proteins of the present invention.

Nucleic acid molecules that hybridize to polynucleotides of the invention can be isolated, e.g., from genomic or cDNA libraries. In order to identify and isolate such nucleic acid molecules the polynucleotides of the invention or parts of these polynucleotides or the reverse complements of these polynucleotides can be used, for example by means of hybridization according to conventional methods (see, e.g., Sambrook (1989), loc. cit.). As a hybridization probe nucleic acid molecules can be used, for example, that have exactly or basically the nucleotide sequence of a part of the sequence shown in SEQ ID No: 1 or sequences complementary thereto. The fragments used as hybridization probe can be synthetic fragments that were produced by means of conventional synthesis methods and the sequence of which basically corresponds to the sequence of a nucleic acid molecule of the invention.

Preferably, the nucleic acid molecule of the invention is DNA or RNA.

An alternative embodiment of the invention relates to a vector comprising an above defined nucleotide acid molecule.

The vector of the present invention may be, e.g., a plasmid, phagemid, cosmid, fosmid, BAC, virus, bacteriophage or another vector used e.g. conventionally in genetic engineering, and may comprise further genes such as marker genes which allow for the selection of said vector in a suitable hosts and under suitable conditions.

Furthermore, the vector of the present invention may, in addition to the nucleic acid molecule of the invention, comprise expression control elements, allowing proper expression of the coding regions in suitable hosts. Such control elements are known to the artisan and may include a promoter, a splice cassette, translation initiation codon, translation and insertion site for introducing an insert into the vector. Preferably, the nucleic acid molecule of the invention is operably linked to said expression control sequences allowing expression in eukaryotic or prokaryotic cells.

Many suitable vectors are known to those skilled in molecular biology, the choice of which would depend on the function desired and include plasmids, phagemid, cosmids, fosmid, BAC, virus, bacteriophages and other vectors used conventionally

in genetic engineering. Methods which are well known to those skilled in the art can
5 be used to construct various plasmids and vectors; see, for example, the techniques described in Sambrook (1989) loc. cit. and Ausubel (1998) loc. cit.. Alternatively, the nucleic acid molecule and vectors of the invention can be reconstituted into liposomes for delivery to target cells. Thus, according to the invention relevant sequences can be transferred into expression vectors where expression of a
10 particular (poly)peptide/protein is required. Typical cloning vectors include pBscpt sk, pGEM, pUC9, pBR322 and pGBT9. Typical expression vectors include pTRE, pCAL-n-EK, pESP-1, pOP13CAT. Typical prokaryotic cloning and expression vectors include: plasmid vectors like the pUC series (e.g. pUC18, pUC19), pGEM series (e.g. pGEM 7zf+, Promega, USA), pET series (e.g. pET22B, Novagen, USA), pBBC
15 1MCS series, pNOF (GL Biotech Germany), pCR-TOPO series and pCR Blunt (Invitrogen, USA), pBluescript series, pCAL series and pBC series (Stratagene, USA); Fosmid vectors like pEpifos5 and pCC1 (Epicentre, USA); Cosmid vectors like the Expand series (Expand I, II, III) (Roche, Germany), SuperCos (Stratagene, USA), pOJ436; BAC vectors like pBeloBAC, pCC1BAC (Epicentre, USA). However,
20 the present invention also envisages the expression of nucleic acid molecules as disclosed herein in eukaryotic vectors. Preferably, said nucleic acid molecules are linked to "control sequences". Said linking may be direct or indirect and refers, preferably to an operable linkage.

The term "control sequence" refers to regulatory DNA sequences which are
25 necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, control sequences generally include promoter, ribosomal binding site, and terminators. In eukaryotes generally control sequences include promoters, terminators and, in some instances, enhancers, transactivators or transcription
30 factors. The term "control sequence" is intended to include, at a minimum, all components the presence of which are necessary for expression, and may also include additional advantageous components.

The term "operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner.

A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. In case the control sequence is a promoter, it is obvious for a skilled person that double-stranded nucleic acid is preferably used.

Thus, the vector of the invention is preferably an expression vector. An "expression vector" is a construct that can be used to transform a selected host cell and provides for expression of a coding sequence in the selected host. Expression vectors can for instance be cloning vectors, binary vectors or integrating vectors. Expression comprises transcription of the nucleic acid molecule preferably into a translatable mRNA. Regulatory elements ensuring expression in prokaryotes and/or eukaryotic cells are well known to those skilled in the art. In the case of eukaryotic cells they comprise normally promoters ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Possible regulatory elements permitting expression in prokaryotic hosts comprise, e.g., the PL, lac, trp or tac promoter in *E. coli*, and examples of regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (In-vitrogene), pSPORT1 (GIBCO BRL). Typical prokaryotic cloning and expression vectors include: plasmid vectors like the pUC series (e.g. pUC18, pUC19), pGEM series (e.g. pGEM 7zf+, Promega, USA), pET series (e.g. pET22B, Novagen, USA), pBBC 1MCS series, pNOF (GL Biotech Germany), pCR-TOPO series and pCR Blunt (Invitrogen, USA), pBluescript series, pCAL series and pBC series (Stratagene, USA); Fosmid vectors like pEpifos5 and pCC1 (Epicentre, USA); Cosmid vectors like the Expand series (Expand I, II, III) (Roche, Germany), SuperCos (Stratagene, USA), pOJ436; BAC vectors like pBeloBAC, pCC1BAC (Epicentre, USA).

An alternative expression system which could be used to express a cell cycle interacting protein is an insect system. In one such system, *Autographa californica*

nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The coding sequence of a nucleic acid molecule of the invention may be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of said coding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein coat. The recombinant viruses are then used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which the protein of the invention is expressed (Smith, J. Virol. 46 (1983), 584; Engelhard, Proc. Nat. Acad. Sci. USA 91 (1994), 3224-3227).

In plants, promoters commonly used are the polyubiquitin promoter, and the actin promoter for ubiquitous expression. The termination signals usually employed are from the Nopaline Synthase promoter or from the CAMV 35S promoter. A plant translational enhancer often used is the TMV omega sequences, the inclusion of an intron (Intron-1 from the Shrunken gene of maize, for example) has been shown to increase expression levels by up to 100-fold. (Mait, Transgenic Research 6 (1997), 143-156; Ni, Plant Journal 7 (1995), 661-676). Additional regulatory elements may include transcriptional as well as translational enhancers. Advantageously, the above-described vectors of the invention comprises a selectable and/or scorable marker. Selectable marker genes useful for the selection of transformed cells and, e.g., plant tissue and plants are well known to those skilled in the art and comprise, for example, antimetabolite resistance as the basis of selection for dhfr, which confers resistance to methotrexate (Reiss, Plant Physiol. (Life Sci. Adv.) 13 (1994), 143-149); npt, which confers resistance to the aminoglycosides neomycin, kanamycin and paromycin (Herrera-Estrella, EMBO J. 2 (1983), 987-995) and hyg, which confers resistance to hygromycin (Marsh, Gene 32 (1984), 481-485). Additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman, Proc. Natl. Acad. Sci. USA 85 (1988), 8047); mannose-6-phosphate isomerase which allows cells to utilize mannose (WO 94/20627) and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue, 1987, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory

ed.) or deaminase from *Aspergillus terreus* which confers resistance to Blasticidin S (Tamura, Biosci. Biotechnol. Biochem. 59 (1995), 2336-2338).

Useful scorable marker are also known to those skilled in the art and are commercially available. Advantageously, said marker is a gene encoding luciferase (Giacomin, Pl. Sci. 116 (1996), 59-72; Scikantha, J. Bact. 178 (1996), 121), green fluorescent protein (Gerdes, FEBS Lett. 389 (1996), 44-47), β -glucuronidase (Jefferson, EMBO J. 6 (1987), 3901-3907) or secreted alkaline phosphatase (SEAP) (Schlatter et al. (2001) Biotechnol Bioeng. 5, 75(5), 597-606). This embodiment is particularly useful for simple and rapid screening of cells, tissues and organisms containing a vector of the invention.

The present invention furthermore relates to host containing an aforementioned vector or an aforementioned nucleic acid molecule. Said host may be produced by introducing said vector or nucleotide sequence into the host by transfection or transformation wherein the nucleotide sequence and/or the encoded (poly)peptide/protein is foreign to the host. Upon the presence of said vector or nucleotide sequence in the host the expression of a protein encoded by the nucleotide sequence of the invention or comprising a nucleotide sequence or a vector according to the invention is mediated.

By "foreign" it is meant that the nucleotide sequence and/or the encoded (poly)peptide/protein is either heterologous with respect to the host, this means derived from a cell or organism with a different genomic background, or is homologous with respect to the host but located in a different genomic environment than the naturally occurring counterpart of said nucleotide sequence. This means that, if the nucleotide sequence is homologous with respect to the host, it is not located in its natural location in the genome of said host, in particular it is surrounded by different genes. In this case the nucleotide sequence may be either under the control of its own promoter or under the control of a heterologous promoter. The vector or nucleotide sequence according to the invention which is present in the host may either be integrated into the genome of the host or it may be maintained in some form extrachromosomally. In this respect, it is also to be understood that the nucleotide sequence of the invention can be used to restore or create a mutant gene

via homologous recombination.

Moreover, the present invention related to a method for producing a (poly)peptide as encoded by a nucleic acid molecule of the invention, comprising culturing the host of the invention under suitable conditions and isolating said (poly)peptide from the culture.

Isolation and purification of the recombinantly produced proteins and (poly)peptides may be carried out by conventional means including preparative chromatography and affinity and immunological separations involving affinity chromatography with monoclonal or polyclonal antibodies specifically interacting with said proteins/(poly)peptides. Preferably, said antibodies are antibodies of the invention as described herein below.

As used herein, the term „isolated protein“ includes proteins substantially free of other proteins, nucleic acids, lipids, carbohydrates or other materials with which it is naturally associated. Such proteins however not only comprise recombinantly produced proteins but include isolated naturally occurring proteins, synthetically produced proteins, or proteins produced by a combination of these methods. Means for preparing such proteins are well understood in the art. The proteins of the invention are preferably in a substantially purified form. A recombinantly produced version of said proteins, including secreted proteins, can be substantially purified by the one-step method described in Smith and Johnson, 1988.

An alternative embodiment of the invention relates to a (poly)peptide encoded by a nucleic acid molecule of the invention or as obtained by the method of the invention.

Preferably said (poly)peptide or fragment thereof is glycosylated, phosphorylated, amidated and/or myristylated.

Furthermore, the present invention relates to an antibody or an aptamer specifically recognizing the aforementioned (poly)peptide or a fragment or epitope thereof. Said antibody may be a monoclonal or a polyclonal antibody.

The term “fragment thereof” as used herein refers to fragments of said

(poly)peptide/protein which are characterized by their capability to induce an immunological response in an immunized organism. Said response may be induced by the protein or fragment thereof either alone or in combination with a hapten, an adjuvant or other compounds known in the art to induce or elicit immunoresponses to a protein or fragment thereof.

The term "epitope" defines a single antigenic determinant. Said determinant is at least a portion of an antigen to which e.g. an antibody specifically binds to by its paratope; see Roitt et. al. (1993) Immunology 3rd edition, Mosby.

A preferred embodiment of the invention relates to an antibody which is a monoclonal antibody.

Said antibody, which is monoclonal antibody, polyclonal antibody, single chain antibody, or fragment thereof that specifically binds said peptide or polypeptide also including bispecific antibody, synthetic antibody, antibody fragment, such as Fab, a F(ab₂), Fv or scFv fragments etc., or a chemically modified derivative of any of these (all comprised by the term "antibody"). Monoclonal antibodies can be prepared, for example, by the techniques as originally described in Köhler and Milstein, Nature 256 (1975), 495, and Galfré, Meth. Enzymol. 73 (1981), 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals with modifications developed by the art. Furthermore, antibodies or fragments thereof to the aforementioned peptides can be obtained by using methods which are described, e.g., in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. When derivatives of said antibodies are obtained by the phage display technique, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the peptide or polypeptide of the invention (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmberg, J. Immunol. Methods 183 (1995), 7-13). The production of chimeric antibodies is described, for example, in WO89/09622. A further source of antibodies to be utilized in accordance with the present invention are so-called xenogenic antibodies. The general principle for the production of xenogenic antibodies such as human antibodies in mice is described in, e.g., WO 91/10741, WO 94/02602, WO 96/34096 and WO 96/33735. Antibodies to be

employed in accordance with the invention or their corresponding immunoglobulin chain(s) can be further modified using conventional techniques known in the art, for example, by using amino acid deletion(s), insertion(s), substitution(s), addition(s),

and/or recombination(s) and/or any other modification(s) known in the art either
5 alone or in combination. Methods for introducing such modifications in the DNA sequence underlying the amino acid sequence of an immunoglobulin chain are well known to the person skilled in the art; see, e.g., Sambrook (1989), loc. cit..

The term "monoclonal" or "polyclonal antibody" (see Harlow and Lane, (1988), loc. cit.) also relates to derivatives of said antibodies which retain or essentially retain
10 their binding specificity. Whereas particularly preferred embodiments of said derivatives are specified further herein below, other preferred derivatives of such antibodies are chimeric antibodies comprising, for example, a mouse or rat variable region and a human constant region.

The term "scFv fragment" (single-chain Fv fragment) is well understood in the art
15 and preferred due to its small size and the possibility to recombinantly produce such fragments.

The term "specifically binds" in connection with the antibody used in accordance with the present invention means that the antibody etc. does not or essentially does not cross-react with (poly)peptides of similar structures. Cross-reactivity of a panel of
20 antibodies etc. under investigation may be tested, for example, by assessing binding of said panel of antibodies etc. under conventional conditions (see, e.g., Harlow and Lane, (1988), loc. cit.) to the (poly)peptide of interest as well as to a number of more or less (structurally and/or functionally) closely related (poly)peptides. Only those
25 antibodies that bind to the (poly)peptide/protein of interest but do not or do not essentially bind to any of the other (poly)peptides which are preferably expressed by the same organism/tissue as the (poly)peptide of interest, e.g. by a crenarchaeote, are considered specific for the (poly)peptide/protein of interest and selected for further studies in accordance with the method of the invention.

30 In a further alternative embodiment the present invention relates to a transgenic non-human mammal whose somatic and germ cells comprise at least one gene encoding a functional polypeptide selected from the group consisting of:

- (a) the polypeptide of the invention;
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution.

A method for the production of a transgenic non-human animal, for example transgenic mouse, comprises introduction of the aforementioned polynucleotide or targeting vector into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with the invention in a method for identification of compounds, described herein below. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., *Gene Targeting, A Practical Approach* (1993), Oxford University Press. The DNA of the embryonal membranes of embryos can be analyzed using, e.g., Southern blots with an appropriate probe; see supra. A general method for making transgenic non-human animals is described in the art, see for example WO 94/24274. For making transgenic non-human organisms (which include homologously targeted non-human animals), embryonal stem cells (ES cells) are preferred. Murine ES cells, such as AB-1 line grown on mitotically inactive SNL76/7 cell feeder layers (McMahon and Bradley, *Cell* 62:1073-1085 (1990)) essentially as described (Robertson, E. J. (1987) in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*. E. J. Robertson, ed. (Oxford: IRL Press), p. 71-112) may be used for homologous gene targeting. Other suitable ES lines include, but are not limited to, the E14 line (Hooper et al., *Nature* 326:292-295 (1987)), the D3 line (Doetschman et al., *J. Embryol. Exp. Morph.* 87:27-45 (1985)), the CCE line (Robertson et al., *Nature* 323:445-448 (1986)), the AK-7 line (Zhuang et al., *Cell* 77:875-884 (1994)). The success of generating a mouse line from ES cells bearing a specific targeted mutation depends on the pluripotency of the ES cells (i. e., their ability, once injected into a host developing embryo, such as a blastocyst or morula, to participate in embryogenesis and contribute to the germ cells of the resulting animal). The blastocysts containing the injected ES cells are allowed to develop in

the uteri of pseudopregnant non-human females and are born, e.g. as chimeric mice. The resultant transgenic mice are chimeric for cells having either the recombinase or reporter loci and are backcrossed and screened for the presence of the correctly targeted transgene (s) by PCR or Southern blot analysis on tail biopsy DNA of
 5 offspring so as to identify transgenic mice heterozygous for either the recombinase or reporter locus/loci.

The transgenic non-human animals may, for example, be transgenic mice, rats, hamsters, dogs, monkeys, rabbits, pigs, or cows. Preferably, said transgenic non-human animal is a mouse.

10 The figures show:

Figure 1A shows the Structure of humic acids (after Stevenson 1982).

Figure 1B shows the Structure of fulvic acids (after Buffle 1977).

15 Figure 2 shows the Structure of polyvinylpyrrolidone (Monomer)

Figure 3 shows an example of an 2-phase PVP-low-melting agarosegel for simultaneous DNA purification and size resolution (Discontinuous Affinity-Gelelectrophoresis; DAG). The PVP-agarose-phase (the first phase) typically takes
 20 up approximately 1/4 to 1/3 of the gel but may take up maximally 80-95 % of the gel. After migrating through the PVP-phase into the agarose-only phase gel segments containing DNA fragments of interesting size may be excised and appropriately treated. The DNA migrates from minus (-) to plus (+).

25 Figure 4 shows an example of an 2-Phase Column chromatography for size resolution and affinity purification of DNA. In the first phase DNA in solution is resolved from inhibitors by size-exclusion chromatography. After passing this part of the composite column, the DNA passes a phase containing an affinity matrix (e.g.
 30 PVPP) to selectively bind and retard inhibitors. The DNA elutes from the column largest molecules first. The liquid flow is driven by hydrostatic or peristaltic pressure.

Figure 5 shows a pulsed-field gelelectrophoretic (PFGE-) separation of metagenomic DNA isolated from soil (method A). The high molecular weight DNA is concentrated in a compression zone above 600 kbp. Yeast genomic DNA and a commercial size marker are added for reference.

Figure 6 shows an insert analysis of metagenomic clones containing soil-derived DNA. Clones were digested with Not I.

Figure 7 shows an expression screening of arrayed fosmid clones containing metagenomic, soil-derived DNA. The encircled clone shows a halo of substrate degradation indicating hydrolase enzyme activity.

Figure 8 shows the quantification of metagenomic soil DNA in fractions eluting from a column after PVPP/sepharose 2B chromatography.

The DNA bands of fractions 6 - 22 were quantified by fluorescence intensity comparison after gel electrophoresis in a 1% agarose gel containing ethidium bromide. Gel documentation and analysis was done using GeneTools software from SynGene (UK). A 1% agarose gel was chosen to concentrate heterogenous DNA fragments in a single band for the sake of simplifying quantification. This was achieved yet at the price of size-resolution. The apparent comigration of eluting DNA with the 23 kbp marker band therefore is an underestimation of true maximum DNA fragment sizes. Lane 1: λ Hind III Marker DNA

Figure 9 shows the separation of soil metagenome DNA from humic substances by chromatography on a PVPP/sepharose 2 B column.

Spectral absorption of eluting fractions 1–22 at 260 (DNA and humic acids) and 230 nm (humic acids) are plotted along their ratio and the relative DNA amounts as determined by agarose gel electrophoresis (see figure 8). A high A260/A230 ratio indicates pure DNA with low humic/fulvic acid contamination as can be seen in fractions 8 and 9. Absorption at 260 nm and 230 nm rises in two peaks in later fractions (12 and 17) possibly due to two size-populations of humic acids.

Figure 10: Restriction digest (*NotI*) of randomly selected environmental fosmid clones separated by pulse field gel electrophoresis. Lanes 1,2: DNA size standards. A band of 7.5 kb visible in all lanes corresponds to the fosmid vector.

- 5 Figure 11: Phylogenetic tree based on full-length 23S rDNA sequences of Bacteria, Archaea and of sequences obtained from marine environmental genomic clones. Different alignment filters were used to evaluate the phylogenetic reconstruction. The tree topology shown here is based on a maximum parsimony analysis (using 1048 conserved positions selected by a positional variability filter). Closed circles indicate
- 10 branching points supported by different phylogenetic methods and filters in reconstructions with the 23S and the corresponding 16S rDNA tree (see Experimental Procedures in appended example 5).

- Figure 12: Schematic representation of the archaeal fosmid clone 29i4/SEQ ID NO:
- 15 1. Different shadings indicate the phylogenetic affinity of the putative protein-coding genes to archaea (diag. stripes), bacteria (dots), bacteria and archaea (vertical stripes), or archaea, bacteria and eukarya (grey). Hypothetical genes with no homologs are shown without fillings. ORF numbers match those in Table 1.

- 20 Figure 13: Phylogenetic analysis with selected sequences of the ETF-like protein family. Homologs of FixA proteins from archaea form a monophyletic group with the FixA proteins of nitrogen fixing bacteria and of *Thermotoga maritima*, clearly distinguished from "housekeeping" ETF proteins (Etf β homologs) of bacteria, archaea and eukaryotes. A third distinct subgroup (termed FixA paralog) is formed
- 25 by a few as yet uncharacterized sequences from bacteria and archaea. For details of the phylogenetic reconstruction see Experimental Procedures in appended example 5.

30 Examples

The following examples illustrate the invention. These examples should not be construed as limiting: the examples are included for purposes of illustration and the present invention is limited only by the claims.

5 **Example 1 Generation of a fosmid library from soil metagenomic DNA**

1.1 Preparation of High Molecular Weight DNA from Soil (A)

Soil was collected from the upper layer of a partially ruderalized sandy ecosystem in Weiterstadt (Germany). About 50 g were suspended in 300 ml of buffer (pH 8, 20 mM Tris-HCl, 10 mM ϵ -aminocaproic acid, 10 mM EDTA) and incubated at 4°C for
0 15 h with gentle shaking. The suspension was sieved to remove larger particles and after centrifugation of the filtrate (5000 x g 30 min.) the resulting microbial fraction was embedded in agarose plugs (0.5 % low-melting SEAPlaque, FMC Bioproducts). The plugs were incubated at 37°C for 1 hour in lysozyme buffer (100 mM EDTA pH 8.0, 10 mM Tris-HCl pH 8.0, 50 mM NaCl, 0.2% Na-deoxycholate, 1% laurylsarcosin,
5 2 mg/ml lysozyme), then transferred into ESP solution containing 2 mg/ml proteinase K, 1% lauroyl Sarcosin and 0.5 M EDTA and incubated at 50°C for 24 h under gentle rotation and with 1 exchange of ESP buffer. 3 agarose plugs were placed in a 1% agarose gel (Sigma A-2929) that contained 2 % PVP (Sigma PVP-360) in the upper part and no PVP in the lower part. Pulsed-field electrophoresis was performed in a
0 CHEF-DR II PFGE machine (BioRad) at 10°C, 6 V cm⁻¹ for 20 h with 1 to 4 s pulses (Figure 5). A slice of agarose containing DNA in the size range of >30 kbp was cut out of the gel inserted into appropriately sized slots in a second gel and re-electrophoresed for a second size selection. After electroelution the resulting DNA was dialyzed and concentrated in a microconcentrator (Vivascience).

5 **1.2 Preparation of High Molecular Weight DNA from Soil (B)**

Alternatively total soil DNA was extracted using a protocol modified from Zhou and coworkers (Zhou et al., (1996) loc. cit.). Soil (5 g) was resuspended in 13.5 ml extractionbuffer (100 mM Tris/HCl pH8.0, 100 mM Na-EDTA, 100 mM Na-phosphate pH 8.0, 1.5 M NaCl, 1% CTAB) followed by an optional 3 cycles of
0 freezing in liquid nitrogen and boiling in a microwave oven. After adding 1.5 ml of lysozyme solution (50 mg/ml) and 30 min. incubation at 37°C, 200 μ l proteinase K

solution (10 mg/ml) were added followed by another 30 min. incubation at 37°C. Then 3 ml 10 % SDS was added followed by 2 hours incubation at 65°C. After centrifugation (10 min., 6000 x g, room temperature) the supernatant was collected and the pellet reextracted twice for 10 min. at 65°C with 4.5 ml of extraction buffer and 1 ml of 10% SDS. All supernatants were united and extracted with an equal volume of chloroform/isoamylalcohol (24:1 vol/vol). DNA was precipitated from the aqueous phase with 0.6 vol isopropanol, pelleted by centrifugation (16000 x g, 20 min., room temperature), washed in 70% ethanol and dissolved in 200µl TE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA). Depending on the soil sample this solution was yellowish to dark brown. This DNA routinely defied any enzymatic manipulation by restriction enzymes and could be successfully used for PCR only at dilutions below 1/1000. It was therefore electrophoresed either in a constant voltage electrophoresis using a 0.5 % agarose 2-phase gel with 2 % PVP in the first phase (4 hours at 60 V) or in a 2-phase PFGE gel as described in (A).

Example 2: Library construction and analysis

About 0.5 µg of purified DNA was enzymatically treated to prepare 5' phosphorylated blunt ends and was ligated to the linearized and dephosphorylated fosmid vector pEpiFOS-5 (pEpiFOS Fosmid Library Production Kit, Epicentre). After in vitro packaging into lambda phages (Epicentre) and infection of E.coli strain EPI100 (Epicentre), cells were plated on LB medium containing 12.5 µg/ml chloramphenicol. The colonies were transferred to individual wells of 384-well microtitre plates containing 50 µl of LB with 12.5 µg/ml chloramphenicol and 7 % glycerol (v/v) and were incubated at 37°C for 24 hours. The library was stored at -80°C.

After blunt-end cloning into the fosmid vector pEpiFOS-5 about 50000 colonies were obtained per µg of soil DNA. Our final library was constructed from a single ligation mixture and contained 25278 clones arrayed in 66 384-well microtitre plates.

Restriction analysis of 30 randomly chosen clones with NotI showed insert sizes between 32.5 and 43.5 kbp, with an average of 36.5 kbp which corresponds well with the insert size range acceptable for this type of vector (Figure 6). The total library therefore contained an estimated 0.9 Gbp of environmental genomic soil DNA, which represents 225 genome equivalents assuming a 4 Mbp average genome size. Most

inserts analyzed exhibited complex patterns after NotI digestion, suggesting that these clones contained DNA with high GC content.

Example 3: Screening of the library

5 The arrayed clones of the library were plated

a) onto LB-agar to grow the cells for a subsequent preparation of fosmid-pools as a resource for sequence-based screenings (using PCR and degenerate primers to generate metagenome sequence tags e.g. as probes for hybridisation) and

10 b) onto LB-agar containing specific substrates for the detection of enzymatically active colonies e.g. through scoring of clearing zones around the colonies (Figure 7) and

c) onto LB-agar for growth and subsequent overlay with a lawn of indicator organisms to score for recombinants producing antibiotics.

15 Example 4: Purification of soil metagenomic DNA by use of a 2-phase gel-permeation/affinity column

Metagenome DNA from soil was extracted by gentle chemical lysis as described before (Zhou et al., (1996) *loc. cit.*). Crude DNA extract was passed over a 2-phase column by gravity-flow using a borosilicate glass column (BIORAD #737-0717). The
20 column (7x150 mm) was packed with a lower phase of 1ml PVPP (SIGMA # P-6755) and an upper layer of Sepharose 2B (SIGMA # 2B-300) to a final volume of 5 ml. After equilibrating the column with 20ml of running buffer (100 mM NaCl, 10 mM Tris, 1 mM EDTA; pH 8.0) separation of DNA from humic and fulvic acids was
25 initiated by applying crude DNA extract (ideally 1-5 % resin volume) to the top of the column. Eluting fractions of 300 µl were collected dropwise (6 drops) and subsequently analysed. Relative DNA content was quantified by comparing the fluorescence of DNA bands in each fraction after electrophoresis on a 1% agarose gel containing ethidium bromide (figure 8) and documentation using a CCD camera system (GeneGenius) and GeneTools software package (both Syngene, UK). The
30 spectral absorption of fractions at 230 nm and 260 nm was measured spectrophotometrically.

Fractions 8 and 9 contain pure clonable DNA. Analysis of later eluting fractions 10–12 show humic acid contamination as can be seen in the decrease in the ratio OD260/OD230. This is due to the significant rise of the OD230 values - the maximum absorption for humic acids. As humic substances also absorb at longer wavelengths the OD260 values increase similarly. The separation of fractions containing pure DNA and fractions containing humic substances can be improved further by adjusting the ratio of the loaded volume of crude DNA extract to the resin volume of the column in a way known to those skilled in the art and by adjusting the DNA content of the loaded samples.

Example 5: Exemplified identification of genes of a microorganism as part of a metagenome in accordance with the invention.

Traditionally, soil microbiology has focused on the description of cultivable microorganisms, while functional aspects of soil microbial communities have mainly been restricted to bulk studies that involved the monitoring of substrates and levels of end products. The application of molecular techniques to microbial ecology revealed that many of the microbial transformations in the environments might be performed by organisms that have not yet been cultivated and thus far remained uncharacterized (Pace (1997) *Science* 276, 734-740, Hugenholtz et. al. (1998) *J. Bacteriol.* 180:4765-74). Soil was confirmed to be particularly rich in microbial diversity based on phylogenetic studies with 16S rRNA genes directly amplified from environmental samples (see e.g. Hugenholtz et. al. (1998) loc. cit., Borneman et al. (1996) *Appl. Environm. Microbiol.* 62, 1935-1943., Barns et. al. (1999) *Appl. Environm. Microbiol.* 65, 1731-1737, Dunbar et. al. (1999) *Appl. Environm. Microbiol.* 65, 1662-1669). Frequently, evidence was even found for the existence of microorganisms from divisions that were not previously associated with soil habitats and of which no cultivated relatives. Among one of the most striking discoveries was the frequent detection of non-thermophilic members of the archaeal kingdom Crenarchaeota (DeLong (1998) *Curr. Opin. Genet. Dev.* 8, 649-654). 16S rDNA sequences of these archaea were first identified in marine picoplankton (DeLong (1992) *Proc. Natl. Acad. Sci. U S A* 89, 5685-9, Fuhrman et. al. (1992) *Nature* 356, 148-9) and then found in freshwater habitats (Hershberger et. al. (1996) *Nature* 384,

420, Schleper et. al. (1997a) *Appl. Env. Microbiol.* 63, 321-323, McGregor et. al. (1997) *Appl. Env. Microbiol.* 63, 1178-1181, Jurgens et. al. (2000) *FEMS Microbiol. Ecol.* 34, 45-56) and in soils from various locations in the United States (Bintrim, et. al. (1997) *Proc. Natl. Acad. Sci. USA* 94, 277-282, Buckley et. al. (1998) *Appl. Environ. Microbiol.* 64, 4333-4339), Finland (Jurgens et. al. (1997) *Appl. Environ. Microbiol.* 63, 803-805), Japan (Ueda et. al. (1995) *Eur. J. Soil Sci.* 46, 415-421, Kudo et. al. (1997) *Biosci. Biotechnol. Biochem.* 61, 917-20) and Germany (Sanda, et. al. (1999) *Appl. Environ. Microbiol.* 65, 3293-3297, Ochsenreiter et. al., in preparation). The ubiquitous ecological distribution of crenarchaeota was very surprising, because their cultivated relatives are exclusively (hyper)thermophiles isolated from terrestrial and marine hot springs.

Quantitative estimates have demonstrated the significant occurrence of non-thermophilic crenarchaeota in marine habitats (Massana et. al. (1997) *Appl. Environ. Microbiol.* 63, 50-6, Karner et. al. (2001) *Nature* 409, 507-510), in freshwater sediments (McGregor et. al. (1997) loc. cit.) and in soil (14,19). Some crenarchaeotal lineages were shown to be specifically associated with plant roots, indicating that the organisms might play a role in the ecology of the rhizosphere (Simon et. al. (2000) *Environ. Microbiol.* 2, 495-505, Chelius & Triplett (2001) *Microb. Ecol.* 41,252-263). While 16S rRNA studies have provided insights into the huge extent of microbial diversity novel approaches are being sought to be able to functionally characterize those microorganisms that have escaped classical cultivation approaches.

Inspired by the rapid advances in microbial genomics of cultivated organisms, a novel approach has recently been initiated to characterize uncultivated organisms that have solely been predicted in rRNA gene surveys. It involves the construction of complex habitat-specific gene libraries by direct cloning of genomic fragments from environmental samples into cloning vectors (DeLong (2001) *Curr. Opin. Microbiol.* 4, 290-295). With the help of phylogenetically relevant gene markers, such as e.g. rDNA genes, large genomic fragments of specific phylotypes can be isolated from these libraries. Yet at present the full potential of this approach cannot be realized as technical constraints severely hamper the cloning of large DNA inserts, particularly from microbial consortia of inhibitor-rich environments like soils or sediments. The approach has successfully been applied to characterize uncultivated, marine

microorganisms: Several genome fragments of the symbiotic crenarchaeote *Cenarchaeum symbiosum* and of marine archaea representing abundant components of the picoplankton in North Pacific and Antarctic waters were retrieved

from BAC and fosmid libraries (Stein et. al. (1996) *J. Bacteriol.* 178,591-599,

5 Schleper et. al. (1998) *J. Bacteriol.* 180, 5003-5009, Béjà et. al. (2000a) *Environ.*

Microbiol. 2, 516-529, Beja et. al. (2002a) *Nature* 415:630-633, Béjà et. al. (2002b)

Appl. Environ. Microbiol. 68, 335-45). A comparison of crenarchaeotal fosmids

revealed significant genomic divergence even in clones with identical 16S rRNA sequences (Béjà et. al. (2002b) loc. cit.). The diversity of large photosynthetic gene

10 clusters of proteobacteria was analyzed from marine planktonic genomic samples

(Beja et. al. (2002a) loc. cit.). A novel type of rhodopsin, termed proteorhodopsin that

functions as a light-driven proton pump was discovered in the genomic fragment of

an uncultivated marine γ -proteobacterium (Béjà et. al. (2000b) *Science* 289, 1902-

1906). The analyses of large genomic regions of hitherto uncultivated organisms

15 also provided the basis for functional studies, including the monitoring of protein

activities in the environment (Beja et. al. (2002a) loc. cit., Beja, et. al. (2001) *Nature*

411, 786-9) and the characterization of proteins after expression in the surrogate

host *E.coli* (Béjà et. al. (2000b) loc. cit., Schleper et. al. (1997b) *J. Bacteriol.* 179,

7803-7811).

20 The colocalization of functional, metabolic genes with phylogenetically ascribable

genetic markers like rRNA genes provides insights into the physiological potential of

uncultivated microorganisms. Clearly the likelihood of a physical colocalization of

such markers on a contiguous cloned DNA stretch will be directly linked to DNA

fragment size. This highlights the relevance of cloning large uninterrupted DNA

25 fragments which is technically very difficult to achieve particularly from microbial

consortia of inhibitor-rich environments like soils or sediments. We have developed

procedures for the efficient purification of large DNA fragments by eliminating the

polyphenolic compounds that heavily contaminate soil DNA. We have constructed

complex genomic libraries and used these to isolate fragments from non-

30 thermophilic crenarchaeota. While direct cloning of large DNA from soil samples has

been demonstrated earlier (Rondon et. al. (2000) *Appl. Environ. Microbiol.* 66, 2541-

2547), our study represents the first genomic characterization of a lineage of soil

microorganisms that has solely been predicted by PCR-based studies.

5.1 Experimental Procedures

Preparation of DNA from Soil

Soil was collected from the upper layer (0 to 5 cm) of a partially ruderalized sandy ecosystem ("Am Rotböhl") near Darmstadt (Germany) in early Spring 2001. DNA was prepared as described in Example 1.1, supra. About 50 g were suspended in 300 ml of buffer (20 mM Tris pH 8, 10 mM ϵ -aminocaproic acid, 10 mM EDTA) and incubated at 4°C for 15h with gentle shaking. The sample was filtered to remove larger particles, and the microbial fraction was centrifuged and embedded into agarose plugs (0.5% low-melting SEAPlaque, FMC Bioproducts). These plugs were incubated at 37°C for 1 hour in 100 mM EDTA, 10 mM Tris-HCl pH 8, 50 mM NaCl, 0.2% deoxycholate, 1% lauroyl sarcosine, 1 mg/ml lysozyme, then transferred into ESP buffer (2 mg/ml proteinase K, 1% lauroyl sarcosine, 0.5 M EDTA) and incubated at 50°C for 24 h with gentle rotation and with one exchange of buffer. Agarose plugs were placed in a 1% agarose gel (Sigma A-2929) which contained 2% polyvinylpyrrolidone (VP-360, Sigma) in the first half and no PVP in the second half. Pulse field gel electrophoresis was performed at 10°C, 6 V cm⁻¹ for 20 h with 1 to 4 sec pulses in a CHEF-DR II (BioRad). DNA of > 30 kb was extracted from the gel and submitted to a second size selection using a regular agarose gel. After electroelution the resulting DNA was dialyzed and concentrated in a microconcentrator.

Library construction

Purified DNA (0.5 μ g) was enzymatically treated to prepare 5' phosphorylated blunt ends and was ligated into fosmid vector pEpiFOS-5 (pEpiFOSTM, Epicentre). After *in vitro* packaging into lambda phages, the infected cells were plated on LB⁺ medium (containing 12.5 μ g/ml chloramphenicol). The colonies were transferred to 384-well microtitre plates containing 50 μ l of LB⁺ medium and 7 % glycerol (v/v). The plates were incubated at 37°C for 24 hours.

16S rDNA diversity studies

Primers specific for the domain Archaea (20F/958R, DeLong (1992) *Proc Natl Acad Sci USA* 89: 5685-9) and Bacteria (27F/1391R, Reysenbach and Pace (1995) *Archaea: a laboratory manual* (Cold spring Harbor Laboratory Press)) were used to

amplify 16S rDNA fragments from the DNA used for constructing the large-insert library. The fragments were subsequently cloned into pGEM-T-easy (Promega) and sequenced. The ARB-software package (Ludwig et al. (1998) *Electrophoresis* 19: 554-568.) was used for alignments and phylogenetic analyses of the partial 16S rDNA genes.

Screening and sequence analysis of Fosmid clone 29i4

Plasmid DNA from the library was prepared from pools of 384 clones and screened by PCR with archaea-specific 16S rDNA primers (DeLong (1992) loc. cit.). A product of correct size (950 bp) was obtained from pool 29. It was randomly labelled with digoxigenin (Roche Biochemicals) and used as a probe in colony hybridization to identify the individual clone (i4). A subclone library was prepared from the fosmid DNA by mechanical shearing and cloning of 2-3 kbp fragments into pGEM-T-easy (Promega). The ends of the cloned DNA fragments were sequenced with vector primers using ABI3700 capillary sequencers. Remaining sequence gaps were closed by primer walking with sequence-derived oligonucleotides.

Sequence annotation

The ORF identification and automatic gene annotation were done with the help of the MAGPIE program package (Gaasterland and Sensen (1996) *Biochimie* 78: 302-310). The Wisconsin Package (Heidelberg Unix Sequence Analysis Server, HUSAR) was used for additional searches with GCGBLAST, identification of PFAM domains and transmembrane segments, for secondary structure prediction and peptide motifs. A tRNA gene was identified using the tRNA scan server (<http://www.genetics.wustl.edu/eddy/tRNAscan-SE/>). Multiple alignments were done with PILEUP and CLUSTAL and manually corrected in SEQLAB. The sequence was deposited in EMBL under the accession no. AJ496176.

Phylogenetic Analyses

The ARB-software package (Ludwig et al. (1998) loc. cit.) was used for alignments and phylogenetic analyses of full-length 16S and 23S rDNA genes from Archaea and from the marine environmental clones of a euryarchaeote 37F11 (Béjà et al. (2000a) loc. cit.), and of two crenarchaeota (4B7, Stein et al. (1996) loc. cit., Béjà et al. (2002b) loc. cit.). The topologies of the 23S rDNA tree were evaluated using the

maximum parsimony (parsimony interactive) and the distance matrix (Felsenstein correction) method with different alignment filters (gap-filter, positional variability filter, maximum frequency filter). The topologies of the corresponding 16S rDNA tree were evaluated using the same methods and in addition the maximum likelihood (fastDNAmI) method with different alignment filters as described above. Phylogenetic analysis of the putative FixA gene (ORF12/SEQ ID NO: 24) was performed using the protein parsimony (PROTPARS) and neighbor-joining (NEIGHBOR) programs from PHYLIP version 3.6 and PAUP version 3.1.1. The same overall topology was found with both methods based on 211 conserved positions from a sequence alignment of 33 FixA/ETF β (flavoprotein containing electron transport chain) homologs.

5.2 Results

Construction of a fosmid library from soil DNA

For preparation of high molecular weight DNA biomass from a soil sample was embedded into agarose plugs prior to lysis. The resulting DNA appeared heavily contaminated with polyphenolic compounds (i.e. humic and fulvic acids) as indicated by a dark brownish appearance. In the process of optimizing the subsequent purification steps, a pulse field electrophoresis procedure was developed that involved an agarose gel with two phases. It allowed purification of the DNA from soil substances through a PVP (polyvinylpyrrolidone) containing phase. In a second phase the PVP was subsequently eliminated, while a first size selection of the DNA was achieved. Highly concentrated, pure and clonable DNA in the size range of 30 to 100 kb was recovered in this one-step electrophoresis procedure, thereby minimizing shearing effects that tend to occur in repeated electrophoresis procedures. The approach has been successfully applied to different soil samples, like ruderal, agricultural and forest soils, for rapid preparation of pure and concentrated high molecular weight DNA (data not shown). After blunt-end cloning into the fosmid vector about 50,000 colonies were obtained per μ g of soil DNA. Our final library contained 25,278 clones.

Restriction analysis of 30 randomly chosen clones using *NotI* showed insert sizes between 32.5 and 43.5 kb. The library therefore contained approximately 0.9 Gbp of environmental genomic soil DNA. 27 of 30 inserts analyzed exhibited complex

patterns after *NotI* digestion, suggesting that these clones contained DNA with high G+C content (*NotI* recognition sequence: GCGGCCGC; Fig. 10).

Sequencing of insert ends from 2688 clones revealed in about 25 % of the sequences significant similarities to protein genes from the data bases (e-values of <

5 10^{-10} in blastx searches). Among these were homologs of proteins from lineages that are typically found in soils, i.e. streptomycetes, clostridia and bacilli (data not shown). However most of the sequences did not show significant similarities to known protein genes. Together, these results confirmed that a great diversity of genomic DNAs was contained in the library.

10 To further monitor the diversity of the DNA used for construction of the library, a PCR-based 16S rDNA survey was performed using a primer set specific for Bacteria. 16S rRNA gene fragments affiliated with eight different bacterial phyla were identified in a random sample of 50 different sequences, many of which are typical for soil microbial assemblages, i.e. Actinobacteria, Chloroflexi, α, β, ϵ Proteobacteria,

15 Planctomycetes, Acidobacterium/Holophaga, Cytophaga.

Identification and analysis of a genomic clone from non-thermophilic crenarchaeota in soil.

Using a multiplex PCR approach and 16S rDNA-specific probes, an archaeal fosmid clone was identified in the library. The insert of clone 29i4/SEQ ID NO: 1 was entirely

20 sequenced and comprised 33,925 bp with an average G+C content of 40 %. It encoded a complete 16S and 23S ribosomal RNA operon, one tRNA gene and 17 predicted protein-encoding genes. The 16S RNA gene was 97 % identical over 711 positions (*E.coli* positions 8-719) to sequences previously recovered in a PCR study from the same soil (Ochsenreiter and Schleper, manuscript in preparation) and 95-

25 97 % identical to sequences obtained from a soil in Wisconsin (Bintrim et al. (1997) loc. cit.). The ribosomal RNA operon of clone 29i4 consisted only of the 16S and 23S rRNA genes, without linked 5S rRNA or tRNA genes similar to cultivated thermophilic and uncultivated marine crenarchaeota. Phylogenetic analyses with the complete 16S rRNA and 23S rRNA genes confirmed the affiliation of clone 29i4 with

30 the crenarchaeota (Fig. 11). As predicted by phylogenetic analyses with partial 16S rRNA sequences, it formed a sister group to the uncultivated marine organisms. The phylogenetic tree in Fig. 11 is based on complete 23S rRNA genes from known

archaeal genomes and from environmental genomic fragments of marine archaea. The same branching orders were found in 16S rRNA phylogenies (see legend to Fig 11).

Ten of the 17 predicted protein-encoding genes showed significant similarity to genes of known function, two were conserved hypothetical genes, five open reading frames did not show any similarity to sequences in the databases (Febr. 2002). Eight of the predicted proteins showed highest similarities to archaeal homologs (Table 1 and Fig. 12). A family B DNA polymerase shared 46 % identical positions with its closest homolog from *Cenarchaeum symbiosum* (Schleper et al. (1997) J Bacteriol. 179, 7803-11). Although the C-terminal end of about 90 amino acids was not encoded on fosmid 29i4, all conserved exonuclease and polymerase motifs typically found in this class of DNA polymerases were identified in the deduced amino acid sequence.

Two other predicted proteins belonging to conserved archaeal protein families were an asparagine synthetase and a phosphoserine phosphatase, both involved in amino acid metabolism. Two putative glycosyl transferases (ORF07/SEQ ID NO.: 14 and 10/SEQ ID NO.: 20) shared significant similarities with homologs from the crenarchaeote *Sulfolobus solfataricus* and from the euryarchaeote *Pyrococcus* ssp., respectively. In contrast, a putative polyhydroxyalkanoate synthase (ORF04/SEQ ID NO.: 8) and a second α/β -hydrolase (ORF02/SEQ ID NO.: 4) were most closely related to bacterial proteins.

A gene cluster was identified with high similarity in structure and sequence to the fixABCX operons found in many symbiotic nitrogen-fixing soil bacteria. Based on its similarity to the components of the flavoprotein-containing electron transport chain (ETF) that is involved in β -oxidation of fatty acids in mitochondria and some bacteria, the operon was proposed to encode a flavoprotein-containing electron transport chain (Weidenhaupt et al. (1996) Arch Microbiol 165: 169-178). In symbiotic bacteria the operon is co-regulated with other genes involved in nitrogen fixation (Gubler and Hennecke (1998) J Bacteriol 170: 1205-1214). FixABCX genes have also been identified in the genomes of several other bacteria and some thermophilic and hyperthermophilic archaea, i.e. in *Sulfolobus solfataricus*, *Thermoplasma acidophilum*, *Pyrobaculum aerophilum* and *Aeropyrum pernix*. Phylogenetic analyses

of the putative FixA gene from 29i4 and homologs from completely sequenced microbial genomes indicated a close affiliation of FixA from 29i4 with other archaeal proteins. Together with FixA proteins from nitrogen fixing bacteria they formed a distinct subgroup within the Etf β /FixA superfamily (Fig. 13).

- 5 A sensory histidine kinase was identified in close proximity but oriented in opposite direction to the fixABCX operon on clone 29i4. While the C-terminal half of the protein exhibited the conserved motifs typically found in sensory histidine kinases, no similarities to known proteins that might give hints to its specific role in sensing were found in the 350 amino-acid long N-terminal part.
- 10 Using PCR primers targeting the ends of the insert of clone 29i4 and internal protein coding genes, no contiguous genomic fragments overlapping clone 29i4 could thus far be detected in the library. However, additional archaeal clones were identified from non-thermophilic crenarchaeota in a second library that contained another 1.5 Gbp of DNA from the same soil sample. One of these clones was identified with
- 15 archaea-specific 16S rRNA probes (as used for clone 29i4) and two other clones were identified through sequencing of insert ends from 768 randomly chosen clones. The sequence analysis of these genomic fragments is under way.

5.2 Discussion

- 20 The direct cloning of high-molecular weight DNA from soil is particularly difficult due to the occurrence of polyphenolic compounds that co-purify with DNA and severely inhibit PCR amplification reactions, hydrolysis by restriction enzymes, ligation and cloning procedures (Trevors and Van Elsas (1995) *Nucleic Acids in the Environment: Methods and Applications*, Springer Verlag, Berlin; Young et al. (1993) loc. cit.). Therefore, any purification protocol for DNA from soil samples must remove the
- 25 phenolic compounds. Protocols have been developed that involve the addition of hexadecylmethylammoniumbromide (CTAB, Zhou et al. (1996) loc. cit.) or polyvinylpyrrolidone (PVP, Trevors and Van Elsas (1995) loc. cit.) in the extraction buffers, because these compounds complex polyphenolics or reduce their electrophoretic mobility when included in electrophoresis procedures (Young et al.
- 30 (1993) loc. cit.). However, such compounds in turn have to be efficiently eliminated (e.g. by extraction, electrophoresis or affinity chromatography) because they inhibit enzymatic treatments of the isolated nucleic acids. Due to these difficulties most

purification procedures either result in high quality DNA suitable for PCR amplification of gene fragments but too highly degraded for cloning of large fragments or it results in high molecular weight DNA that is not pure enough for cloning procedures. (refs above and own observations). Therefore, the device of the

5 invention was developed for the electrophoresis procedure described herein above (two phases in which the DNA is first purified from polyphenolics in a PVP containing phase and subsequently cleaned in a second phase, thereby minimizing shearing effects that occur in repeated electrophoresis procedures). This technique allows reproducible obtainment of highly concentrated and pure, high molecular weight
10 DNA. While purification techniques similar to those described by Zhou et al. (1996) loc. cit.) that was used by Rondon et al. (2000) loc. cit.) for cloning large DNA fragments were not applicable to many of our samples, the novel PVP electrophoresis procedure yielded reliable results with different soil samples of varying organic and humic acid content. Successful purification of the DNA was
15 independent of the lysis procedures that we used, i.e. direct lysis of soil samples as in Zhou et al. (1996) loc. cit.) or lysis of microbial fractions as described here.

The complex environmental libraries constructed exemplarily by using the device and method of the invention contain a large fraction of the total genomic content of a soil microbial population, which has been referred to as the soil "metagenome" (Rondon
20 et al. (2000) loc. cit.). The library characterized here was constructed in a BAC-derived fosmid containing cos-sites for *in vitro* packaging with lambda phages. Said vector yielded significantly larger clone numbers than using classical BAC-vectors (data not shown) and it allowed the direct cloning of undigested DNA by blunt-end ligation, thereby avoiding any bias introduced by restriction digests. Using 50 g of
25 soil, a library with 0.9 Gbp of environmental DNA was constructed which represents approximately 225 genome equivalents assuming a 4 Mbp average genome size. Using archaea-specific 16S rDNA probes the fosmid clone 29i4 was identified. Sequence analysis demonstrated that a contiguous genomic fragment of non-thermophilic soil crenarchaeota was isolated: (i) Phylogenetic analyses based on the
30 complete ribosomal 16S and 23S RNA genes indicated the specific affiliation with the crenarchaeotal clade as predicted in PCR-based studies. (ii) genes affiliated with archaea were found dispersed over the entire clone insert (iii) G+C content and

codon usage of the predicted protein genes were similar over the entire insert and (iv) the deduced amino acid sequence of a DNA polymerase gene showed greatest similarity to its homolog from the uncultivated marine symbiont *Cenarchaeum*

symbiosum. Functional and biochemical analysis of the latter protein had confirmed the predicted non-thermophilic phenotype of this crenarchaeote (Schleper et al. 1997) J Bacteriol. 179, 7803-11).

On the other hand, significant differences to the content and structure of genomic fragments from uncultivated non-thermophilic marine archaea revealed that crenarchaeota from soil have significantly diverged from their relatives in other environments. An unusually large gap in the 16S-23S RNA operon and the lack of a GSAT gene (glutamate semialdehyde aminotransferase), which was consistently found to be directly linked to the ribosomal operon on all marine crenarchaeotal genome fragments (Béjà et al. (2002) loc. cit.) indicates the difference. Direct comparison of the soil fosmid clone to the genomic clones obtained from marine planctonic crenarchaeota and from the symbiont *C. symbiosum* revealed only one related protein encoding gene, i.e. the putative DNA polymerase. Genes on clone 29i4/SEQ ID NO: 1 appeared to be less densely packed than in genomes of other archaea. Only 69 % of the sequence encoded RNA or protein genes. There was the large intergenic region in the 16S/23S rRNA cluster of 830 bp, which is atypical for ribosomal operons in crenarchaeota. Another large non-coding region of 2787 bp was found between ORF10/SEQ ID NO: 20 and fixA. No apparent genes or distinctive structural features, e.g. repetitive elements were identified in the non-coding regions.

The genomic information contained on fosmid 29i4/SEQ ID NO: 1 gives first insights into metabolic properties of crenarchaeota from soil and can serve as a basis for functional genomic studies. Beside genes encoding proteins for "house-keeping" functions (replication, amino acid metabolism), two α/β type hydrolases so far seem to be particularly found in soil crenarchaeota. One of them encodes a putative protein involved in the synthesis of polyhydroxyalkanoates. The operon encoding FixABCX revealed a putative flavoprotein containing electron transport chain that is commonly found in symbiotic nitrogen-fixing bacteria. A detailed phylogenetic analysis indicated that the putative FixA protein of 29i4 is most closely affiliated with

archaeal FixA homologs and not with the FixA proteins of Bacteria or the paralogous ETF proteins from other species of Archaea, Eucarya or Bacteria. Therefore, it seems unlikely that the fixABCX genes of 29i4 have been acquired by horizontal gene transfer e.g. from symbiotic nitrogen-fixing bacteria which might reside in the same soil habitat. They rather seem to originate from a common ancestor of the Archaea. None of the obligately aerobic archaea that contain the fixABCX genes is known to be capable of fixing nitrogen. Expression analysis of this operon in well-studied thermophilic model organisms, such as *Sulfolobus solfataricus* might shed light on its physiological role in crenarchaeota.

Table 1:

Predicted RNA and protein encoding genes in the archaeal fosmid 29i4.

ORF	Nt range	protein size	putative function	most similar ortholog*	Phyl. aff. ^a	Comments
01	1-2367	789 aa (C-term. truncated)	Family B DNA polymerase pfam00136	AAC62689 Cenarchaeum symbiosum (0.0)	AEB	
RNA	3548-5018		16S RNA	85% Identity to 16S RNA of Cenarchaeum symbiosum		
RNA	5847-8818		23S RNA	77% Identity to 23S RNA of Cenarchaeum symbiosum		
02	8968-9849	293 aa	α/β hydrolase pfam00561	AAD02150 Pseudomonas stutzeri (e -17)	AEB	Catalytic triade Ser/Asp/His; close homologs are from bacteria
03	9888-10205	105 aa	Hypothetical	none		
04	10219-11304	361 aa	PHA Synthase (α/β hydrolase, pfam00561)	P45366 Thiocystis violacea (e -66)	B	No homolog in archaea but PHB production has been described in Halobacteriaceae
05	11285-11866	193 aa	Hypothetical	none		
RNA	12414-12502		tRNA ^{Asp}			
06	13017-13454	145 aa	Hypothetical	none		
07	14324-15238	304 aa	Glycosyl transferases group 1, pfam00534	AAK41834 Sulfolobus solfataricus (e -11)	AEB	Transfer of ADP, UDP, GDP, CMP linked sugars
08	15716-17407	563 aa	Asparagine synthetase pfam 00310	AAB99117 Methanococcus janaschii (e -67)	AEB	
09	17492-18157	221 aa	Phosphoserine phosphatase pfam00702	AAB86099 Methanothermobacter thermautotrophicus (e -14)	AEB	
10	18377-19588	403 aa	Conserved hypothetical	CAB50138 Pyrococcus abyssi (e -91)	A	Homologs only found in P. abyssi, P. horikoshii, A. pernix, Glycosyltransferase group 2 domain; pfam 00535
11	20630-21793	387 aa	Transmembrane protein	BAB50489 Mesorhizobium loti (e -31)	B	Domain pfam 01173
12	24580-25461	293 aa	Fix A pfam01012	P53578 Azotobacter vinelandii (e -23)	AB	Paralogs of ETF β : electron transfer flavoprotein β subunit
13	25458-26741	427 aa	Fix B pfam00766	P53578 Clostridium saccharobutylicum (e -45)	AB	Paralogs of ETF α : electron transfer flavoprotein α subunit
14	26738-28615	625 aa	Fix CX	NP_454687 Salmonella enterica (e -36) FixX: Thermoplasma volcanium (e -08)	AB	Fused protein of FixC and FixX
15	29228-31465	745 aa	Sensory transduction histidine kinase, pfam00512	BAB73503 Nostoc sp. PCC 7120 (e -12)	AEB	
16	32505-33023	172 aa	Hypothetical protein	none		
17	32918-33925	335 aa (truncated)	hypothetical	none		

5

*proteins are designated by their gene identification numbers followed by the species name. The e-values of blastx searches are added in brackets.

^a Phyl. aff. = Phylogenetic affinity, denotes occurrence of homologs in Archaea (A), Bacteria (B) or Eucarya (E).

SEQUENCE LISTING

5 <110> BRAIN AG

10

<120> Isolation and Cloning of DNA from Uncultivated Organisms

15

<130> G1184 EP

20

<170> PatentIn version 3.1

25

<210> 1

<211> 33925

30

<212> DNA

<213> Crenarchaeote

35

<400> 1

cacagccttg ttgatcataa cattaaaact taattcttcc agagaaattc tttttttctc 60

caagttttct gcaattgatt gcaactatctt ttttatcttt tcctttgctc tttcaaagtc 120

40

tttttctgaa aatattttctt taaggacatt taatatgtca tagaaagctt gtcttattat 180

	tggaggtgta tgagacttct tccctgttaa tcctttaaca tctacagttc cgtcctccaa	240
	tacacctagg taattttttt ttagttcact aaaaaccacg tagcgatata ttttatctat	300
5	ctccaaatct atgcctagtt cttttttaga ccaggatgaa attccactta atccttcctt	360
	ggaaggattc ttttaggaaca gagaatccgt atcaccgtaa ataacctcaa tcttttcttc	420
10	gttgcathtt tcaatagttt ttgttgtggt catccttcca accgctgcgg tagcctcagc	480
	tacaggtaaa caatagagcg gaaatatttc agcaccata accccatacg tagcatttaa	540
	aataaccttt atggcctgac tgataacact gtatagctgt ttatcctctt tatccaaaga	600
15	attatccttt gatagatatt tgtaataatt aacccttaga tcccttaggg ttccatatcaa	660
	tatggaggtc atcccttgct tttccttgca aacccaatgg tttgtttgct caatatgtgt	720
20	tgatggatcc cttctgcaat tttcatgagg acaattgact gtttcgtaag ataaattgtg	780
	aactttaatt atgctaggat acagactagc aaaatctaca actatgacat tgaaatgaat	840
	tcctaaaacg ggctcaacca ccagacctcc tcgatatttt ttttccttta taatggcaac	900
25	tgtagacgat gttccttttt tctgtaattc atctttacgg ggaataatga tattttgctg	960
	cctatgttca aaaaacatca tggaccta at ccattgattc accccgaatc ttgttatatc	1020
30	ttctattgac attcgggata tcctagaaat gataatcaac aattttatca gtaaattgtc	1080
	attgaaagat gtcagacgaa atgtcaagtc tgcattcttg aggcaatact cggccagttt	1140
	ttccaatgga agatcaccta tgctttcatc aaagtctatt ttgactcgt ttaatagggc	1200
35	ttcgcagata gcatttaaag taaactcaga gtatttatga ctaaaagcat aattctgtac	1260
	agatttatht tgaaatgtcc tgaataaatc gatatggatt ccatgcttta aggaaaccgg	1320
40	atccgcctga atacccttt ttataaaaga gtctttttta actaaaatag gcaccaattc	1380
	tttactaatg ggttttttgt gtacagggtc tatcgatggg tottgagatc tagcatataa	1440

	ataaggtaaa tcaaaatcat caccattaata ggttaaaact attggataat tttgaataat	1500
	agcaaaaact tttagtatca tgtctttttc gctatcacat aattcaacgg ttgttgaatc	1560
5	tagtttagag ggatcaaaat ttggatcttt tcttaagaag aatacctttc taaatccatc	1620
	cgatgccgat aaaccactg cagtaattac tttatcgtga tctctggctg tgggcatcct	1680
10	tccctcttca gagtccactt ctatatcaa agaaattctt ttgatatccg gaataggctg	1740
	gtttaacaat cttgaccact ttattagaaa ctcatatat tcaactctt ttggctcatt	1800
	ttctttaaaa ttaggtttta tgagattatc cagatattcg tcaactttct ctggcattgg	1860
15	gaattcatga aaaactaaat tattgcctat tctgttataa aacgctcccg gaattagacc	1920
	caaatcgaat aaataacttt catggtattt gatatcagcc tccaagaag taaccttttc	1980
20	cctaaaacta ctatctgttc caccaatgga aaggggggtca ggggcaatta ttttaaaaac	2040
	cgatatttcc ttgtcctcaa tatcgtccat tttttttatt ttttctagtc taaacctatg	2100
	tggctctttg ctaactattg ttttaacctg atcagaatag agttccttta caaagcaata	2160
25	aggttgatgt ttattttatat gattttcaat aaaagactcg ctccaaaaat atatttgaga	2220
	atcttctgga ttgtaaaact tcaaaaacac tgattttttt tctcctatgt aaacggaaga	2280
30	aagtaacaat gatggaatat tttctggaag ttctttttga taataactat ttttctctc	2340
	cagaggcaag tcggatttta tcaccatatt tttatttcat tcatcctttt aaaaattaat	2400
	gacttgtttt agaaaactat caaatttgat ttacctagca ccatcaaggc tgatgtctga	2460
35	ctaggttatt tttagcatta ctgatttatt tttctttgag gacaaacctt cttaagaata	2520
	taccattat attattcaag cgtgtcaatc gttactcag tattatcaat atcgacagca	2580
40	atgtgagaac aagagggttt ggatgtggat ggatgatcaa tggttgattt tcaaattcct	2640
	gtatgaaata ttgtgagcaa tcgatatatc attagataaa agaagattta gttaggacat	2700

	cttgataata tattcgtgta gattgatttg aaaaaacagc agcaataaga aaatttaata	2760
	cgataaaaata cgataacttc gactaactaa catcccttgg gaaaagaact taatgcccga	2820
5	tccccttttt tttgtttttt aaagaaaaaa gaagatttat attaacaatc ctaccataag	2880
	tagtagaagc cgtccaagac aaaaggcggc gtcggtgatg aagttggttt gtgccataaa	2940
10	gtgatttact gacctccaga tgcgctcggt acattactgg ttaacgatat aatattaaaa	3000
	tagtgggata atggggattc gatcaacaca tcggaactga cttgatggat cggatctgac	3060
	atgggaagat cagacagaca ttgaaaaaga ttcattataa cagcacaaaa gatatcattg	3120
15	caggcattgt ttgtatgtgt gtgacaatac aatacagcat atgottgtga gttaatctaa	3180
	caaataacca aataaacaaa tcagaagggt attagagttt ttctttcttt ttcgatcgta	3240
20	ctctctttcc ctgcttttta aaggcaggga gaaaactcaa taaacgttct tttgtgtttg	3300
	ccaatggctt ttccctcttt ctcttaccgt cttctcggat gtgagggcgg aggcgggaag	3360
	gttggtggca gagaccaaag caacgcgtat atacaccata aagcaaaagt caaccgatag	3420
25	gtaacaaatg gcgcacgttt gtgttttttt cottgtggcg ttttgcctct ctcaaaaaag	3480
	gcaaggcaaa acccatatgt gtgcgtttgt catctgttat gtttttcacc atcatcattt	3540
30	ttttttgaat ccggttgatc ctgccggacc cgactgctat cagagtggga ctaagccatg	3600
	cgagtcaaca tagcaatatg tggcatacgg ctacagtaaca cgtagtcaac atgcccaggg	3660
	gacgtggata acctcgggaa actgaggata aaccgcgata agtcactact tctggaatgg	3720
35	gtaatgactt aaatctatat ggccccctgga ttggactgcg gccgatcagg ctgttggtga	3780
	ggtaatggcc caccaaact gtaaccggtg cgggctctga gaggaggagc ccggagatgg	3840
40	gcactgagac aaggggcccag gccctatggg gcgcagcagg cgcgaaacct ctgcaatagg	3900
	cgaaagcctg acagggttac tctgagtgat ttccgttaag gagatctttt ggcacctcta	3960

	aaaatggtgc agaataaggg gtgggcaagt ctggtgtcag ccgccgcggt aataccagca	4020
5	ccccgagtgg tcgggacgtt tattgggcct aaagcatccg tagccggttc-tacaagtctt	4080
	ccgttaaate cacctgctta acagatgggc tgcggaagat actatggagc taggaggcgg	4140
	gagaggcaag cggactcga tgggtagggg taaaatccgt tgatccattg aagaccacca	4200
10	gtggcgaagg cggcttgcca gaacgcgctc gacggtgagg gatgaaagct gggggagcaa	4260
	accggattag ataccgggt agtcccagct gtaaacgatg cagactcggg gatgaattgg	4320
15	cttcattgcca attcagtgcc gcagggaagc cgttaagttt gccgcctggg gactacggtc	4380
	gcaagactga aacttaaagg aattggcggg ggagcaccac aaggggtgaa gcctgcgggt	4440
	caattggagt caacgccgga aatcttaccg ggggcgacag cagaatgaag gtcaagccga	4500
20	agactttacc agacaagctg agaggaggtg catggccgct gccagctcgt gccgtgaggt	4560
	gtcctgttaa gtcaggtaac gagcgagacc cctgcctcta gttgctacca ttattctcag	4620
25	gagtagtga gctaattaga gggactgccg tcgctgagac ggaggaagga gggggctacg	4680
	gcaggtcagt atgccccgaa accctcgggc cacacgcggg ctgcaatggt aaggacaatg	4740
	agtatcgatt ccgaaaggag gaggcaatct ctaaacctta ccacagttat gattgagggc	4800
30	tgaaactcgc cctcatgaat atggaatccc tagtaaccgc gtgtcactat cgcgcgggtga	4860
	atacgtccct gtccttgca cacaccgcc gtcgcttcac cgaagtgggt tcttggcgag	4920
35	gtgatgccta attggtacta tcgaacctgg ggtcagcaac gagggagaag tcgtaacaag	4980
	gtggccgtag gggaacctgc ggccggatca cctccttagt tatcatatct tgcaacacag	5040
	aacaaaatag aaaaaagag aaaaatgggt gggaatgaag gaaaaactct acccaccgtt	5100
40	taatttggtt ccttgggatc ttggtcagct tggtttaca acatgaatgc tgcagagtat	5160
	acatcacaca tgcaaaaaca aagcctgcag tggatatctg gcaagtgtta taatggacat	5220

	ggatagggat atgggcatgg atgtgggcaa cacaacacaa agtagtggtg ctagaccaga	5280
	tccgctctgc tcagtgaagag tggaacaaat ttgctagtga cctgtctcta tctgaatgaa	5340
5	tgtgtctgtc tgtctgggtc tttgcgtatg cgtaccogtc ccgtgccaaa gagtcgggtc	5400
	gtgggtaagc caacacaata accatcagag taaaaaaaaac agagtaatgc acgcacacac	5460
10	acatgtgcac acagaaaaag cagaaaggaa aagagaaaga aagaaaagga aggaaagaaa	5520
	gaaaaaaaaagt gaatacggat gcaattcttt gtcactaaac tgaggagtgt gagagacaaa	5580
15	ggatgaaggt gacatagcag gcaacaccta gacaaatata agaaggttgt tgtttgcgat	5640
	gatgtcgtgt ataggatcaa accattaaaa gatataacaa tacattagat atattaaaaa	5700
	tataatgatt cacgtaaaaa tgaaatagtg aaaaattata aaaaaatgta tatctgggtg	5760
20	tagtgcagtg taatacataa aaagcgaatg attttatcat gaaaaccgaa aattaattag	5820
	ttattaccat caatagaaaa caaaattgga aactggaact ggtgcaaaga cgccggttgg	5880
	tggatgactc ggcttgataa gcgaagaagg acgtggcaag ctgcgataag cctggggtag	5940
25	gtgcatgca cgcgcgatcc cgggatgtcc gaatgaggtc tctctttaca ctcccttgct	6000
	ttgttgcttg ggagagcgaa ccgtccgaag tgaagcatct gagtaggacg aggaggagaa	6060
30	atcaattgag attccgtcag tagcggcgag cgaaagcgga acagcccaa ctgaatctgc	6120
	cgtggtaaca cggcagagat gtggtgttgc gggttatagc cataggatcc tgcctttgga	6180
	gctgaagtgt actggaatgt accggaacag agggtgatac ccccgtaggc aaatggaggc	6240
35	gggattctgc tatatccaga gtagctggcc ttggcagtgg ccagtgaagg tgggtgaaag	6300
	tagtatccaa ggctaaatat tcatcaagac ctagagaaaa ctagtaccgt gagggaaagt	6360
40	tgaaaagtac cccggaaggg ggggttaaaag cgcctgaaac caaccggtta cagacgtgta	6420
	tggctcgaaa ggataaaatc tagagtcata cgttccgtct agaaacacgg gccagggaga	6480

ttgctgtcat ggcaagctta accttttaca aagggaatgc gaagggaaac cgaatttgcg 6540
 ctttttctct ttattgagaa aagaggcaat ggatctgaaa gggctctcaag tcatggcagt 6600
 5 aaggctagaa accggacgat ctattcctgg ataagacgaa ggtgagtga aactcgctgg 6660
 aggtctgcaa gggtcctgac gtgcaaatcg gtcccctgat ctgggattag gggcaaaaa 6720
 10 ccaatctagt ccggtgatcg ctagttccca ccgaagtga tgcagtcct gccttagctg 6780
 agatggcctg tattgtagag caccgatcgg gcgtaaggg ctgaaagag ctgccatcc 6840
 attcgaactc cgaatatacg ggcgtcgtag aagctaggag gcgggtttat gtggggtaag 6900
 15 cctcataacc gagaggggga caaccagac taaagttaag gtccccaat gtctactaag 6960
 tgtcaaacca aagggtgttt tcgagcagag acagcaggaa ggtaggctca gaagcagcca 7020
 20 ttcttttaaag agtgcgtaac agctcacctg ccgagctcga aagccccgaa aatgtacggg 7080
 gctcaagtag actaccgata ctttagacca ccgacgatgt cggcgctgg taggtgggcg 7140
 tagtgtttgg gtagaagctg ggctgtaaag tccagtggac cgaactacta gtgcagatcc 7200
 25 tgggtgtagt aacagcatag ccgggtgaga atctcggcga ccgcatgggc aagggtttcc 7260
 cggcaatgcg tcatcagccg ggagttagcc ggtcctaaaa acaacctcaa cagaattgtt 7320
 30 tgaatgggaa actggttaat attccagtgc cttgaaagt cgttaacacc ttttctgtcg 7380
 cttccggata gggtaagcag aaccgtcgtt ctgtccaagt attctagctt tgaggagtac 7440
 cgtaatggcg agaataaaag cgagatacga atggcccttc gcaaggaggg tttgcttgag 7500
 35 tccaggagac actgaaagca gaaacaggga gatactttca agaccgtacc gagatccgac 7560
 actggtgccc tggatgagaa gtctaaggtc tatcgggtat accgtatggc aagggaactc 7620
 40 ggcaaaatag ctccgtacct atggtataag gaggcctgc agtttttacg aggagtaggg 7680
 attgcaggtc gcagtgacta ggggggtccc actgtttaat aaaaacacag gtggctcgcta 7740

gtccgaaagg atgtgtatgg cctctgtatc ctggccagtg gcggtaccta aaacctgggt 7800

acaaccgggc taagggccgc taaacgccgg gagtaactct gactctotta aggtagccaa 7860

5

atgccttgtc gggtaagttc cgacgtgcat gaatggaaca acgagggcc cgctgtccct 7920

gcctacaacc cgtgaagcc acataacgtg gacgaacagt ccacgaacct ctgtcgggga 7980

10

gagaagaccc tgtggagctt tactgcagcc tgttggtgag atatggttgc aaatgcagag 8040

agtagctggg agccgttatg gtcagttctc cgggactgat ctaggcgaca gtgtaacacc 8100

15

agccatttgt taccgtatcg ctaacctgct tatgcagga catcggcagg tgggcagttc 8160

ggctggggcg gcacccctt gaaaatgtat cgagggggcc caaagattgg ctcaggcggg 8220

20

acagaactcc gccggtgagg gcaaagccaa aagccagtct gactggattc ccaatgatac 8280

gggattcaga ggcgaaagcc gggcttagcg atccatcatg tctcactat tgggggctgg 8340

tggtgacaga aaagttaccc tagggataac aggctcgtcg cgggcgagag ctcccatcga 8400

25

ccccgcggtt tggtacctcg atgtcggctc ttcccatcct ggttctgcag caggagccaa 8460

gggtggggct gctcgcccat taaaggggaa cgtgagctgg gtttagaccg tcgtgagaca 8520

30

ggtcggtctc tgcctgacag gggcgtggtt gtctgagggg aagttgcccc tagtacgaga 8580

ggaacagggc agcgcagcct ctggtttatc agttgtccga cagggaagc tgagcagcta 8640

agctgttttag gataactcct gaaagcatct aaggaggaag ctttcccga gacaagacaa 8700

35

ccttcgtaa ggagaagggc ggccatagaa gatggcggtt atggaatgga ggtgtaagca 8760

ccaagctttc aagcgaggtg ttcagcctgc catcaccaat agcccaacgc acctgttgac 8820

40

aaaacaaaa aaaccgacag acagaaaaaa ttgaaaatct ataactaaat atacatattt 8880

ttttgttggg tcattatttc atgcgtaaag agtcaattat agaccaattt gatatatcta 8940

	ctgattattg ttatatagaa ttttttaatg gatattgato ataaaatttt agtatatttc	9000
	atattatcta ttaacaaaat aattattaca atggggttgg tttcggatag acaaagaaac	9060
5	gagacaatgg attttataaa aatactggga tataacatca gatataataa aatagatcaa	9120
	gtcaagtcaa atgaaacat aattctgctt catggtatag gagcttccgc agaacgatgg	9180
	tcagaattag tcccatTTTT gtataattgc aatataatta taccagacat cattgggttt	9240
10	ggttacagtg aaaaaccaag gatagagtac aacatagatt tatttgtaaa gtttttggat	9300
	gaattgtttc tgaaacttga aatcaaaaac cccataataa tgggttcgct ttttggtggt	9360
15	caattgattt tagaatatta tttcaggcac aaagactttt ttaaaaaaat gattctagt	9420
	tccccggccg gtaccaaga gagaccgaca ctagcgtaa ggcaatacac ttactcatgt	9480
	ttatacccaa caagagaaaa taccgaaaga gcatttaaga tgatgtcgca tttcaatcac	9540
20	acagtataag attcaatgat aaaggatttt attaatagaa tgaagcagcc caacgcaaaa	9600
	cactcgtttg tttcaacact tttagcacta aggaaaaata gtgatttaca agacaacctg	9660
25	agggaaatca aaatcccaac tttagtaata tggggaaaag aggacaacac cattccagta	9720
	gaaaatatag agtatttcag gggcatccct tttgtaaaaa catgcataat gagtgattgc	9780
	ggtcattgtc cttttgttga aaagcctctt gagttttata aaatagtcaa agagtttatc	9840
30	gactcctaatt ttctaataata agtattatat tcaacattaa aatattattg aatcaatcca	9900
	cttctatgag taatgagaat gaagaaaata aagatataga ttttaagaaa tccattgaaa	9960
35	aggctgcgga attccagcag gatttggtgc gacagttctc tacaattcaa tacaatgcgt	10020
	ttcagaatat gttttcatct ttgcaaggat ttacaaatta taatgccatg tttaaaacca	10080
	ccgtacagac ggggtggcagg atctcaattc ccgaagcaga aagaaatgct ttggggattg	10140
40	aagaggggtga tctagtccag gttataatta taccgttgac aaggaaaaag aaaaacacaa	10200

gttaaaataa caaatccggt aatgtgtttg aatccatttt ccaatttttg gtaaaacatt 10260
 tttctgtgaa aagttgctag caattagccc tacatgccct gttggaaatt tcatcaggct 10320

 5 tttatcctga cttgaaatta ggttgtttag ggagctactg ctgtcagacg ttacaagggtg 10380
 gtcaaattca gctacaacat taagaacggg aaccttaatg tttgacaaat ttatcttgtt 10440
 10 ttcaccacaca atcatcttgt tttttgcaaa aagggttttg tgatagatat cctttaccca 10500
 ttgcctaaag gtttcccccg caataggagg tgtgtcatac agccatttct ctattcttaa 10560
 aaagttctgt acaaaacttt catcttcaaa gtttttaaataa aaattatagt atttgtttac 10620
 15 accttgcttg aatggtttta gtgatgcata aaccagatac agtaattcat atggaaagtt 10680
 ttctgtgatg gacagtactt tgtcaatata catgtgtctca gccatgtttt ttattacgga 10740
 20 tttgtctttc tcggcatcaa caattggagc aatgggtgact agatttttaa tgtttttttg 10800
 atatagcgaa gtgtacatca aggacattgt acccccatg caatatcctt gtaatgaaat 10860
 ctgatcaatg ttttctatgt tttttatgta ttctacacac tcataaataa acaaattgac 10920
 25 ataatacatca acagtgatgt atttatccag ttttgacggg gggtttccagt caatcagata 10980
 gacatttatg cctgtctcta gcaggttctt tatccaactt ttgtogttct gcagatccaa 11040
 aatatatgat ttgtttatta atgcataaac aatcaacaaa gggtaactga aagtttggtg 11100
 30 ctttaggggt ttataatgta gtaaacggaa gaggggtgtt tcccttatta cctcatattc 11160
 gcttgatcca gtctttatgt tttctatatt cgacaatttt ttctgattt cttttaattt 11220
 35 tgaaatgtta tcaggatccc ttacaaacgt aaaataatca ttactaaat aattaattaa 11280
 ggaattattc atttttttcc tcatttaatt tttttttgat ttccaatgat atttttttga 11340
 tttcatagag attataaaat aacaggtctt tttcttcttt tgacagttgt tgttgtgacc 11400
 40 taaacaaaac ggcatttgaa tcgtaaattt tttgataact tttgatgaca tcaatgctgg 11460

aattcaataa attgttataa ttgattgaaa agtctgttga ctgcaacatt gacgagaaca 11520
catcctcaaa agtatttatg attattttcc taatatcgtc ggggtttttt tcatctatag 11580
5 cagacgatac cttgtttacg gccaacaaat aagcatttat catacgggaa aagtataggt 11640
tgagaaatga ctgatatctc agtaacaaat tacogtgttc ctttaatgta tttaaattaa 11700
10 ggtttgatc attcattaaa gaggtgaatg gccccaaggt tgtaaggcta tttaactctg 11760
ttaattgttt tataaagttt tcaaatacag actgaagtcc tgcctcttca gatagagtgt 11820
ttttactgtt tttttcctct ccaatattgt tattttctaa ttgcaaataa ttgattcacc 11880
15 tcggattcga aactcttgta attgtaataa tttttcatca acaatactat attctttact 11940
ttttgtcatt ataaactttt tttagggtta ggaagaaatg aaattttattg tcatttaaca 12000
20 aacttgaaaa gcaattatcg caggtcaagt gtatgttaca ttgacaattc cttttagtag 12060
agagcagttc ctaaaaactg tcaaaatgga cggatggtaa aaoccatth ccaggcaaaa 12120
gcgcaactgg aatcagaaaag tgttttattcc aaagtttgac catocctctg gactctatth 12180
25 taacgggtct gataatttta atatccttcg gttatacaaa atatggcaca tgacactgat 12240
ttaatgctth tatgctttga ttatcacctt tttatattga aaaaacgact catgcattaa 12300
30 aatagtaaaa ttattctaaa taattttaga tattcacatt taggcaaata attagaagat 12360
aaagccaagg caggcccaac aattttataa ggtaatccta ccaagacgac tgggggtccg 12420
tagctcagca tggatagagc gcctgccttc tatctttacg ggagatagcc ggaagtcgag 12480
35 ggatcgaagc cctccgggcc cgtaaacctt atggggttca aatatttttt cagttctaata 12540
gcccatagaa ggagtatgaa tggacacatc attgcctgaa gttaacaaca accatcacia 12600
40 aaatgaagaa gagaaaggca taattagttg acgaatgtga tgtcggagtt cttttattgt 12660
ttggccggtg tctacgcgtc agaaaactgc caaacgggc tgctcgaggt actggcaaga 12720

catcccgaac agccttgtga tcttctttag ccactgaaaa acatgatggc gtaaggggct 12780

acctcccttc agcgatatat cgctaacegc actgcacttc cctagcggtc agatttcaag 12840

5 tacacagcca tcagtagcct ctaccctgta cgogaatgac gatattcctt gttgcaaact 12900

tttgtttotga atgacatccg tcgcagtatg cagcactagc caggcgtatg catccttcag 12960

10 atcgccaaag caaaagtgga tgtattcggt accccttcca tgtgagcaga cgtccctcac 13020

attgtcactt gcactcctgg gcgtggggcg ctagatttgg gttgccagc aatcattgcg 13080

15 ggcccaaagg cggggtttgt ttctgggtgg ccgggcataa tttgcaaaaa aaggagaaaa 13140

agaacctgac ctttgtcaaa aaatttatat ataggtagag atggttgtct atttttgtat 13200

acaggcacat gagtaattac ccaaatatta tctatggcga aaccatatat agaaacaatg 13260

20 agccaaggca gacatttgcc aaaaccattc gtttgtggat ggattctttc tattgtcccc 13320

cacataacct tcttcccgag gctgatacag catcttagtt tttgggtatg cgcatacccg 13380

cccaccatgt caaaagaatg tatacatacc accgctgggt tggacatgta taaaacatct 13440

25 gaacttgtag gcatgtctca tgcaatgcgc catatggacg atgcggtgac ggtggatttg 13500

ccttactgct gcaaaaactc aaacaaggta ggccccattg gaaggacagc atgttctaag 13560

30 tcggggaaaa cggggcatgc gggtcatttt caggaggaga gtaatatccg tagcccatg 13620

ccaaatgctt gcagtctcaa ggccttgatt tgcgctctgc gcgtttgttt ctttttgaat 13680

acttaactta aggtctggtg cctgttacag ttgatattcc accatagggt gtagatgcag 13740

35 agttattgct attatcagaa gtcggatttc cgggtgtctt gaaatcttgt atggagtgag 13800

tgtgctttgc ggtacatct atcttaacca tataaaaagc agcattgaag acaactgtgg 13860

40 tggaattaat atttgogaat ttatatagc caacaactat ccacgcagga tgagttgaac 13920

taccattggt tgttgtgctg gttatagccc tatagcccct tttatatcat tctatcaagg 13980

tttgtaggat tttgtgaatg gcatatttga atctggctca tcaatatcac aatggcagaa 14040
 attattacta ttgaagacaa tgataggagt ttatgtttat tgccatgcat attaagaact 14100
 5 gaattaaaaa catggcatat aaaattctat ttttgtgact gggataaaga ataacatatt 14160
 ttgtgttatg ggtcatactc aagaattgag actcatcata gttggtgttt ttggttttag 14220
 10 aatcaaaaga agaaaagaac aacagcaata caagttcaaa aagacgttac ccgatcgcta 14280
 atgacaatga tttggaactt ctttctattt ttaactattt tgattagggt acagattcta 14340
 ttgttgtcat tacatTTTTT gccactgtct gccagtttgg gagttggaaa ctgacctttt 14400
 15 ttccttcagt cgcctTTTTTg attatgccat atttatTTtag catttttacg cactctttctg 14460
 caaacagggt ggtttctcca tattctatta gttttatgtt tgtattacca ttttttaagt 14520
 20 atagttcttc aaaaacgggg agtttccagg caacagtggg aacacacgaa actaaggcct 14580
 cagccacagc aatgccaaaa ccctctctgg atgatggaaa aataaagact ttagatttag 14640
 aataaaggct aatcttttct tctcggaga caaagcctct gtgatctata cccgcattac 14700
 25 gtagttttgc agccttatca gggggtatgc gcccaaccat tacaaaatta gattctggtc 14760
 tgagtgtttt tattgcagtc caaatttctt ccagtccatg aaatttttct atccttccga 14820
 30 tacaaagaaa atcaatgtcc tttttattgt tgattactcc tctgttgga tcttttaaaa 14880
 agatattttt atctattcca gttcctacaa tggcaattct gttggtcaga ttttttgcta 14940
 attccctggt ttttttattt gctgtttcct tcaaattatt gatTTtgcta acccctattc 15000
 35 catagactgt gttgagttca tgctttgacg cttgactcac ggtcaaaatc atatcagaat 15060
 ccttaagcat caccgcagta gccttttgta ttagataatt gtacagaaat tcaaaatagt 15120
 40 tcttgcaaat agatatccgc ggttcatggt gatgaaacac aacaaagatc ttggTTTTtg 15180
 gcttgaatag cctaagtagc agccaaagga caatattaga ttcgccccaa gaatccaaaa 15240

tggcaatatac tggacgtttg gcaaggggcc gttaaagcat tgtatgtgtc gggtataatc 15300

ctatttggtat ttttttttgc ggagttatgg ttgtatatct tcaagacaga atagcagccg 15360

5

cttttgtcta tcaggtctgc aacttttttc atccagagaa atcctccggt gtaattcttt 15420

aaacttgaag ggtaaccaa gaataataat ttaatttggt ttgcgttgct ttaatacatc 15480

10 ctggttggtta tatataacta taaagaacac ttattttata attgttattg ttaagaacaa 15540

gataatcaac attttggaca aacatgataa aagaattgaa aaaatgacac aataaattaa 15600

15 aatatttgga taatcgtcat ttacccttca acactgtgga cgatgcaaaa tatcgttttt 15660

gcagtattca tatttttctt tgcgcataac catgaaaatg aggtttggag gttcatcagt 15720

20 tccttggcaa tatgtttttt gaaatattct ccaaatacat ctgtatatgc ggctccaaac 15780

tccagatttg ttcttttca aaaagatgcc catatctttg tgagctgccg ataagctccc 15840

ttttttcaa tttgtccaga taactagaat taaccttgga ttccgtaaaa ccatttttca 15900

25 tggccaaagt attgaggggtg ttgtgtatcc cagaacctat ctgagctgct tcctttatcc 15960

tatacgctat ctcttttga atccctagtt tttctgcaag tttgcggtga accctttttc 16020

30 ctaggttgct atagttattg ccattgtttt gaatattgag tcgcggatct attctcagta 16080

ccgtgtctat cagattagta tctaaaaagg gttcgcgtaa ctctatgctt tgagacatgg 16140

ttattttgtc ctctctttcc agtgtttctt tgtaaagtaa cttaatgtcc tctatcaggt 16200

35 atccctgaat tttttcgtat ccgtgttttt taacaatttt ggaataccag gaatatccgc 16260

caaacagttc gtctgcccc tgacctgtaa gcattaccgc tattccctgt tcgtgagcca 16320

40 atttaaccgc gccatatatt ggaatggcaa cctcaacctg tcccatgttg tcatcttcaa 16380

ttatgctgat tatttttga atggtacttt caacatcact ttcagtcac tgttctattt 16440

ccaacttgag gtcaagtttt tctgctatct caagtgagtt gaggatatca cttgaacctt 16500
 taatcccaga cgtatagcaa ataacttcgg gggccatttg ttttgccaaa tacgctacaa 16560
 5 ttacactgtc aatcccaccg gagaaaacaa taccgatttt tttaaagtca ctacacagtt 16620
 ttctcataga ttcaaccaat gtatcaccat atgcgttaac cgcagaatcg atgtctgtgt 16680
 acaggattga atattttctca catattgatt tttttgtatt tacagaaatc ggaaacaatg 16740
 10 tagtcttgaa attggaggac ctttccttcc gcgaaatgac aagagcatag cctggcaaaa 16800
 gtcttttgat ttggtcggac atagcaattt tccataaggc ttttctttct gatgcaaatg 16860
 15 caatgaaatc actactttca ccatagtaaa tttgtcttac tccaatgcca tcccgtacca 16920
 gcacaatatc tctgtggac tgctctctaa tcgccaaaac ataaattcca tcaagctggg 16980
 taacggttct ctttatagct tcgattagat cgcctttagt gttttgataa tggctcttcaa 17040
 20 gaaggtgaac aataacttca ctatcagtcg aggtagtaaa agtgtgatgt gcagaaaggt 17100
 tctttctgat ttctttatag ttatatattt caccattatg ctccagaatg agttttttat 17160
 25 cacaactcac aaacggctgc tgaccacagg agccaccaac tattgccaaa cgactgtgac 17220
 ctaaaacgtc atgcccctct acctgtgaaa acaatggatt atcaaaggta tcagaataaa 17280
 ctatttgatt ctctgtagac aaacccatgc catcoggacc ccggtttttc atacaggata 17340
 30 gcatttttcc tatcaagggg gcaacatttc tctctttttt acttaaaatt ccaacaattc 17400
 cacacatctt aaaattttcc tatacgggat ttattgatga acaaaatata aaagtaacca 17460
 35 ctattgttgc cattatgggt tctaaacggg tctactctat aaaatcaagg acaccaatca 17520
 tatccggtgt gttacctta ttattgttt catttatata tttatcotta ggcatatatg 17580
 cgatacctat tctgcctgc tttatcatgc acaggtcacc tttagtatcg ccaatagcaa 17640
 40 ttgtattttt tatgtctgca cagattttct ttgcatggat ttccatgtga tatctcttac 17700

acacagaatt cttgcaaaaa cagtctattt tttcccatcc taacggcata tttattttctc 17760

cggtgactat cccattgtct accttcaatt catttgata aaaaaagtcc aaatcaagtt 17820

5 tgttcaccaa ggcctgagca gcaacactgt aactatctgt aattatccct attctgaacc 17880

cttttttctt cagcaaagat atcacctcct ggctgttctt tgcagggggg atggagtcca 17940

10 aagcaatttc tatttccctt tcttctattc ccctaatac agcggctatc ttctgtgtct 18000

taacatagcc tggaatggat ttgtcggact ggatgtgtct gacctgagca tacaagccaa 18060

acttttttga caatacctca attagccttc catcaattag cgtcccatcc atatcaaaaa 18120

15 cggccaatgt agatttaaatt tccataggat acaacaaaca aggaatgtca aagaatatta 18180

ctatttagcg acagcctatt agccaaaatg tttttatagg ttggggacat cattattcaa 18240

20 ttgggatgtc ttgggcacca atttttttat ttcattagat atagccctat aaaaaggtta 18300

cattaaaaag tgttcggttag atcaatttta tgtatgtcat ttataaacga atatgcacat 18360

atagaaatat aaacacatga gattagatta tccacctaac tataccgaga ggataggagc 18420

25 agttagtatc catgcgcttc aaaagattta tgagatcgat tccggaaaga tgcccaagtt 18480

taatggcctg catcagcatc agtctataaa ggccttttgt tatgacgaac tgtcaagcat 18540

30 attccaagaa cttgccatag tcattccagt aaagaacgaa aaaatcagcc ttcttgaagg 18600

agtattgagc ggtattccaa atgaatgtct catcatcata gtttccaata gccaaaggac 18660

tctgtcgac agatttgcca tggagggtga aatggtaagg cagtactcta gttttgcaga 18720

35 caagaaaata atgattattc accaaaatga tcttgagctg gctaatactt ttaagaaaat 18780

aaagtataga tccatcctca acaccaaag tcaggttcgt agtggaagg ctgaaggaaat 18840

gataattgga atattgctgg caaaaatgca cctaaaagag tacattggat ttattgacag 18900

40 tgataattat tttccaggag cagtaaata atagtcaag atctttgcag cgggatttgg 18960

aatggcaacc accccatata gcaatatcag aatatcgtgg cggtccaaac ccaaaatcgt 19020
 aaacaactca ctacaattcc caagatgggg tagaatttca gaatccagta acaaatcct 19080
 5 gaacgctcta atatcccaca tcacagggtt tgaaaggag attatcacga ctggaaatgc 19140
 aggtgagcat gcattatcca tgtcccttgc agaaaatctc aactattcaa gcggatatcc 19200
 gggtgagccc tatgagttta tcaacatttt agaaaagttt ggaggtctac tcccatcaaa 19260
 10 caatcctgac atcatagaaa aggtatcga aatatttcaa atagagacca ggaatccaca 19320
 ctttcatgag gaaaaaggaa atgatcattt ggcaggcatg atgcaagaat ctcttctcgc 19380
 15 aataaacaac agcaaaattt gcaacacaga actgaccagg gaaataaatg accatttact 19440
 catgcttcag gtaaaacaca ataatgatat gaccaaactc aactttaaga aaaaacacct 19500
 20 tataatggat ccataaaaa taatacccat cgacaaatcc gccgaatttg tagttaagaa 19560
 ttctaaaacc ttcatagaa ttggataaaa atatgcagga atgcatattt ttgagaacaa 19620
 caggtttggg aaattttgac tgatttttta gatccctcaa actgcacctt tatccatcct 19680
 25 gttttatcaa gcctgaccaa gcgaatgcat aattatccga ccgtgttttg agcaaccaca 19740
 gaggccactt tttttagaaa caacgtaaag ggataaaaaa cagttgttca ccaacatttc 19800
 30 actagctggt gaataaatta tatcttcaaa cttttattct ccaccctac aaaccgaagg 19860
 atcacagtac tcgcccacg ctacctgaaa aaaaataagc aatagtcagt ttcggatttc 19920
 aaaatttcaa attttccaga gaattaattt tcccctcatc atcaatgccg tcaattactc 19980
 35 tgaagggtat ttttccagct ctttgggtatt ttgttttatt acatattttt ctggatcata 20040
 tccatatttt ttgcctaact cattcatgta ctctgaaca tcatgccaag ctttttgctt 20100
 40 gtcatatgga ttgccattta ggctgctacc tccactgcc aacgtagatt ttgatgaagc 20160
 atccaatgca gtttgggtata gctttgatat cttgctaaat tctcatcag tcaaccttat 20220

atttttatta ttgttggttac tggtcattta ttattacccc attagtaaatt atttgatggt 20280
 caaacttatc tttttctttt gataaaatgg agtcagcctt tgtagcacat tttggatatt 20340
 5
 aaaccaata cgacgcgta cggaagaat aaaagcacct aacacccttc aaaaacattc 20400
 aacgatatga ctgaaagtag ccaaagaatt tgagaatatg ttctttctca tttatcagag 20460
 10 actttttggt tgggtttata attaatgat taacgttctg attgataaaa aagcgcaaaa 20520
 tagcaaacca tgtaaatttg aaaaggggag tacatttggg tatggcttaa caatactgtg 20580
 15 gttgtctcca aaatagtaaa ttttataatc taaaagtaga aaattcccta tgagtgatgc 20640
 tatcgaaaat gtctgatcc ttcagggagg aggatctttg ggtgcatttg gttgcggggg 20700
 ctacaaagca ctagtaaaca ataacataaa acttgatata ctgtctggca catcaattgg 20760
 20 cggtttgaat gccacagtta ttgccggcag taaagaagat cgtccagaaa aatcattgga 20820
 gaatttttgg atggaaatag ctgatactaa taatggtaat attaatatcat accttaattt 20880
 cccctttttt gaaagtcctt ttctgggca aattcctttc cccttggcat cagaatcaac 20940
 25 actatcattc tacagctctg ccatttatgg aaatagaaaa atctttctgc caagatgggg 21000
 acctgaaaat atctttaag atccacagta ttccacacct agcaaatgga catatttgta 21060
 30 tgaccattca ctttggttaa aaaccttgga aaagtacatt gattatagca aattacagcc 21120
 aaacggtaag cccaacgcaa ggctaataat aaccgcagtt aacgtgatga cggcggagcc 21180
 ccttattttt gacagtgcc agcaacaaat aaccccaaaa cacatacttg caaccactgc 21240
 35 ctatccaaca tatttttttc aatgggtgga attggaaaaa gggctttttg cctgggatgg 21300
 aagtttacta agcaataccc cgctaagaga agtaatagac gcatcgcccg caaaggacaa 21360
 40 aagaatcttt ctgtcgaga actatcctaa aaatattgaa aagcttccgt caaacctaca 21420
 ggaagtcaag catagggcaa gagacataat gttcagcgac aagaccgtcc acagtataca 21480

catgtccaaa gcaattaccc ttcaacttaa gcttattgat gatctgtata aaatgctaga 21540

5 gtattacttt aattcagaaa aaatcgagga aaaggagaag ttgaaaaaa ttcgtgcgag 21600

atacaaaaaa gtttcagaag aacacggcgc agagattaaa ggtgtctact atataacacg 21660

ggacgagcca tccccctccc tttatgagaa tgcagacttt tcaaaaaatg caataaaggc 21720

10 atcgattaat gatggagaac aaaaggctga caggataata aaagaaatcc aaacgaaagg 21780

aaaacgaaaa taatgagcca gaaaacacca accaagttgc aatttcaaca accatttttt 21840

15 ttatttggcc tgtattcccc tttttgtcaa aatttttttg caggccaaaa tccaaaccaa 21900

aaggaaaatt cctaattgtct gcaaatttta tttgaaagtg aattatccat attaccatag 21960

agaggcaaat ccagttcgcc ataaaatcct aaacaaaaca atactttttg atccctgcc 22020

20 gaaaagcaac atcagctatc tcagatgtat tggactgagc gctgccatac cacgcgcaac 22080

tttgaaaaca tcgccacacc cataactctt ataccaaatt tttaccaac agaaataaca 22140

catgaattaa aaccaagaa atcagcaata taccattttt gcaaagtcaa ggcatttttag 22200

25 gtaatcgta ataacaaca taacagatta tacagtaaga tcattttggc aggctaaaa 22260

aaagaccgtt ttatttaaga aaaaagcaat tctcgttatg tgggtattat cattgacgat 22320

30 taaccaaatt aaatcaggtc aaatcaggtc aatgactttg cctatttgat aaggtgataa 22380

taccttaggc caaacaatca gcaatatcga ttgttttttg cattaattat ctatttttta 22440

tatttctttt aaaaaacgaa tagaaataat caaatatgtc caaactaaaa tcaagaatta 22500

35 gaaaaatccg tgacagttat ataatatgtt aataataatc aactaatgac aagttcaagt 22560

gaaaatagta gtgataaaga atttgaagag ggcgcagcag gcacaaataa agatagaaaa 22620

40 agtgatccat tgaaagagta tgaaagtaaa gagccaatga caccagcaaa aataaatgaa 22680

ggagaaccaa cggctgtaaa gagagacca tcagaccaa agataacagg agaaggtcaa 22740

acaggagcag ataccgaaca agcagatgaa caattgcgta aacgtggcat gaccaaaatc 22800

~~gattctgatt etttctaacac atctcaataa tcaaaaccaa aataaaaaga ataggatact~~ 22860

5 tcatatcaat atcttttattt tttcttttgt gacgcctttc acccggcaca aattcattat 22920

aaactaatcc aaatgcttcg tttcttgatt gtccaatagc aataaacagc gattagcaaa 22980

10 ccaaacatca acggcatcaa acatcaaaga aataaaaagg tgtgaaaaaa tacataacga 23040

cttgtcgtta tacgaaaatg aaaatatgta ttaaattctt tccataacta gatggaaatc 23100

15 taattttacc ttcaccgata ccattgata agaatcatat ttttctttgc gcattgttat 23160

tattgtgcat agattgatag aaatttacga cgttaaatat ccaggagggt gctatatggt 23220

20 caatttttga atttgaaata ccatgtgctt tatataaaat atttcacgat atatttaaaa 23280

attacaaaaa aacatcgttt gctcgtcatt aaaaacgcaa aaaaacgggc aaaaaaata 23340

ttatatgcat tatatataga atattgtccc tatagtgttt acaatgatac ataattctaa 23400

25 attaacaaca acccttattg ctttgcttat tgttccaata attccaatga tgaccctggg 23460

aataattcca gatgtgattg cacaacagaa cacgacagga attgcagact tgactgaaag 23520

30 caatggcggt ccagatgctg ccgttggcgg cagtagtggc accaacagta gcataggtgg 23580

taacactagt ggctcaagtg aaactatgag tggtataaac ggtggcgaag gcaccgtaga 23640

caaatttcaa tgaggcattt tgccaccct caactttatt aacggccaac cttgccgtct 23700

35 ataagtgatt tgtgacacct tttccttttt attattgtta ccttagttga tatcgaaaag 23760

agactgtatc caaccattta aatatattgg ttattatggc gggttcaggc cctaaaaaag 23820

gaactgcagg caaatagggt tggcctgatg ccaacttttt tggtttgctt caataccaca 23880

40 tctattattt caactccaca aaaacattta cagggatgcc atttgctctc agtccaatcc 23940

taatccttcc ctgctcatat tcaatgtccc caatactatc ttttaatccc cttgaagcaa 24000
 aatgtaaacg attacttttt agctttacct tatatgttcc cgttttttca ttatgggctg 24060
 5 cagacacaaa ctgtatattt tctccgtcaa ccaactcata aagctcccta caaacaggtt 24120
 tgattacctc ccatgtttct ttatctgtta aatccttgtc caataccata aacgggtgct 24180
 10 tctatttttt attgtttaga tatatttgat tatgatatgc agtaaagtat atttctttga 24240
 caaatttaaa atcgatttgc tctatttggg taaattataa aatcatacca acaataagga 24300
 ttttcacaga tgatttgatt ttcacggtga atctgcccag agggatttca aaaggattgc 24360
 15 cagcttatag ttttgtttaa acatttatgt atttaagtag gcataatcag ttagttactg 24420
 tggtttcata cacctaaatt tttgccatat tgtattaaca aaattacggt aaaaaatac 24480
 20 cacgcaataa atcaacttaa atgtaatacc tactttccta tattgccttt ttcaaaacgc 24540
 atttaactcg ttattgagaa ctattaatgt tgaatccaaa tggaacttaa cgcagcagta 24600
 attgtgaaac tcgagccgga tttttctgaa gggaatgtaa gctataattc agacggaaca 24660
 25 cttacagag cagaaacaaa aaacattttg gggccccata gcgcagcagc atccctagca 24720
 gccctgtact caaaagtaaa acatggaacg catgtttctg tgggcacaat gggctctcca 24780
 atagcagaat cggccttaca gcaatctcaa ctgatttgcg acgtgatga actgcatctt 24840
 30 tatagtgatc gcatctttgc aggagccgac accctggcca cagctgaagt tttgatagca 24900
 ggaataaaaa aaatggcaaa tggtaagat gtggacattg ttttctcagg gcacagggca 24960
 35 tctgatggcg aaacagggca aacaggaccc cagacagcat ggaaattagg ttatccgttc 25020
 cttggaaatg ttattgatta cgatattgac gttgtgaaga gaattgtaag ggtacaacgt 25080
 ctaatcaaga tttacggtca tcttgatatt atagaggaga tggaggcgcc tctaccggtt 25140
 40 tttatcacac tggacccatc ctacaatccg tcttttaaca cggtatccca aaggctcaga 25200

ctagcacgaa acctacagga agcccatgat agatcacaaa ggtataagga atatctcaaa 25260

actttcaatg ccatggaact agaagtcaat ccaaagtctg tcggactgcc tggctctccc 25320

5 accatagttt ataaagttga aaaaatacca agggcaaagg caaatagaaa agcagatggt 25380

gtggatgggt ctaaccagga tagtctaagg caggttgac gccgaatcca tgatgtttta 25440

10 gggggtgtag tcataaagtg acatcatcac tatctgcat acctgacgct aaactagacg 25500

aaaggccaaa ccaaatgcc catgttaatg acaaccaga aaaagaaagg ggagacaaca 25560

acaggcatct gtatgttggt atagaacaag aggaaggcac catattacct gtgagttttg 25620

15 aaatgcttgg tgaggcaaga aggctaattg atgattttta tcacaaatac aagccagagg 25680

aaaaagtggg tgcgattata ctggccata acatcaagca cctgtgccag gaactaatcc 25740

20 accatggtgc agacgcagtg atttatgccg accaccgga gtcgccac ccaagaaatc 25800

ttctttatac aaaggttgtc tgccaaattg ctacggacaa agagagcgcc gccagaattt 25860

ggccatcaaa tcccgatttt aacagacccc gttacatgtt tttttccgca gatgacacag 25920

25 gaaggcattt atcatcaacc gttttggcag aattgcaatc agggctggca tcagacataa 25980

acaaacttgt tatcaatgat ttagaaataa ggcataaca caagacaaag ggtaaaccga 26040

30 ttgtctatga aaagacactt gaaatgtaca gaccagactt ttcaggcttt ctttggaaca 26100

ccatactctg cttggataat ataaatcccg agaacagaag gaaattccat ccacaggcat 26160

gcagtataat ccagggcgtc tttcccaaaa tggaaggaga tacggataga aagggtacca 26220

35 taatagagtt cagcccaacc atagcccagg aagaccttag aataaaaata atcaacagaa 26280

gagtaatcaa aagcaaagtc gatttttagca ataaaaaat aatcgtagt tttggaagg 26340

40 gaataaagga gtctcccgaa caaacataa aactgataga gaaccttgca aaggaaatag 26400

aagcagaaat aggaatatca ctgcccattt caaagaaacc ctatccaata agcgaaagtc 26460

tgtcgtcaac ctatatgatt cctgacaggg ttatcggcac aagcggaaga aaggtaaadc 26520
 ctcaggtgta ttttgcaata ggaataagcg gggctgtcca acacatagcc gggatgaaag 26580
 5 aatcggaatt tgtgatttcc atcaatccag acagtgaagc tcccataata gatgaatccg 26640
 atgttttaac caaaggaaaa atcgagcagg tgctgcctct cctgataaat gaattaaaaa 26700
 aatacaaaga gagactgcaa ataccacagg agatagaatg acaatggaaa gttttgatgt 26760
 10 ggcgataatt ggtggagggt ctgctggact tgcggcactt gagcacctct ccaatttggg 26820
 aaaacaggca atcctcatag aggcaggaaa aaaaatagga accaaaaacg tgtctggggg 26880
 15 catattgtat tccaaaaaaa cagcaactgg aaaggtccac aatgtagaag atgtgtttga 26940
 taattttctg gcagacgctc cgctggaaag gaagataata aaatacatgc ttcacgccgt 27000
 ctcaaggga aaagcgttct ctctggacct gactttggca cagcactatc aaacgaattt 27060
 20 tgggtacacc gtctgtctca acaaactact ttcattggtt gcaagggaag catctcaaag 27120
 tgcagaaaaa ctgggtggag ggataataac aggtgtccat ttaagggtoga taatctggaa 27180
 25 agatgacagt accataatta tagagacaga tgaacttgag ccgttccagg taaaggcagt 27240
 cattgcagct gacgggggta actcagaggt tgcgcaaata acaggtgcca gaagcaagtt 27300
 30 cacaccgtct gacctctacc agggcgtaaa ggtggtggca aaattaccag aggggttgct 27360
 tgaagagaga ttcgggggtct cggaacacga gggagcggct cacctttttt caggcgacat 27420
 aacgctaaac cacattggag gagggttcct ttacacaaac agggacacca tctcaattgg 27480
 35 cgcagtatac cattatgact ctctaattga aaagcctaca gagcccaatg cgctgggtcaa 27540
 tgcgttactg tcaaatccgt ttgtgatgga attgataaag gacgaggttc caaggatcaa 27600
 40 ggaggactac agggatcttt caaaggatga agaactaagg attaggttca aatccaataa 27660
 attgataaaa agctggaatg acctacacca cacatattat tcaccatctg ccgttgcaga 27720

gcttgtggcg cagggaaaat acaaatcaag ggaggagatc aaggacaaaa ttgattcatt 27780

gtacaatgag cttgtaacaa aatacaacac agaatttgaa acaaattacg tggagttaga 27840

5 gtacagcgcc aaactgggtc cagatggaaa aaggtgcaga atgaaaaaac cctacttta 27900

aaacatctta tttgtcggg atgctgcggg caggggcatt ttccttgggc cacgcataga 27960

10 gggcctcaac gtaggcattg atgacgcggg tagggccgca gaagctgtct caaagtcaat 28020

agatcaaaat aactttcagt ttgacaacat tgggtgaacgc tacactaaat cagtggatga 28080

15 aagtccatat accgcagaca tgagcaggat cgacgcaaac tatctcaaag ccgttcttga 28140

ttgcacaaaa aaggttccca aaaacactct tgggtttaag tatgggtcta ttgtcaaatt 28200

gatgtcaaat agcaccttta ggaatgtatc cataggaatt gcaaactcta tagggtacaa 28260

20 aaggctttta cctgtgattg agtcagacaa aacctacaat caaattccca tcgagattgc 28320

ggagagaaat ggcaaagatt tgcggaaaag ctattccata gagattccca ccattgccga 28380

gcgtattgct aatctgaact ataatgacga ttcactgtca cacatcaagg ttttgaactc 28440

25 gcaaagtgac tttatgaaaa aaatgggtcca actgtgccct accaaatgct acagtattga 28500

gaatgagcgg ataatgctac agcacgaagg atgcatagag tgtgggacat gcgcaagaga 28560

30 aacagaatgg aggcacctc gtggggaaaa aggaataatc tataattacg ggtaagccat 28620

aaccggaatc catcaacata tcctttctgg aaaaaaagtc ggggataaca cacgcaacaa 28680

aaaaaacaac gaatggattt caggttctaa atttttgggt gtttacacct tatctctgct 28740

35 ttcaccgctt ttattttttt tttgatggat tctatttctt caatcaactc atcaacatac 28800

tccctgattt tttcgtgggt ccccgataac cctccagccc ctgcatactc agttagagaa 28860

40 tcagtttgtt tccggatcac tgtctgcatt ttttcatcaa gctcttcaat caactgtatg 28920

attgacggaa taagcctctc catatgattg gtaaagccac caacaattat aaagcttacg 28980

cagtgtatta atgcgtatta cagtctatat gggtataaac aaccaaaca aatccgaatc 29040
 ---aaaagtaa at gaataacaca taatactaca atggggccatg aaacaaatta catcaaagcg 29100 ---
 5
 tcacatttta agcaacgtca actgctagtt ttgaaagtta tgtattttct tagattat 29160
 tctattctat tttcattggt gtagttgggt gttgcagcag cagttggtgc agttgccaaa 29220
 10
 ctcatgggtca ttttccttca tcgttttttt ctggcatgtc ttttgctggc atatgggaag 29280
 gcaggagac aggtatgaca aatttgaatg tggcgccat ttttccttca ctgccaaagt 29340
 ggtgcagtat ttcacaaatg ccttcgtctt tatctttagt gttgttctg ctgctgctgt 29400
 15
 tgccgccctt gctgtttgtc tctcaaacc atatttttcc accatgctcc tcaacaattt 29460
 tccttgacag gtataggcca aggcgggtc cctgggttga ctttggtgaca aatttctgaa 29520
 20
 acagctgac ccttattttg gagttgagcc caaccccggt gtctgcact gtgactagca 29580
 ccgcgccttc tttctgcctc ccgatgtggt caccaccatt gtcaccacca ccgttgctgc 29640
 tgcgctgct gctatccact ctgccccat tgcctttacc agctgtagca gtgtttgagg 29700
 25
 tatcactttc ctgagaggtg gaagtgaagg gagaagactc accatcact gccgtggaaa 29760
 caacaatctt gccgtcattg gtgaacttca ttgcgttgct cagcagggtg aaaacaacct 29820
 30
 ggcttatctt ttgcggatca cagtctacat aaaaagggtg gttggggcca ttacgggtt 29880
 ctaccactg ctctttttgc tgcgtctctt ttgcgctg ttttgctcct gccgtgcct 29940
 ttacccttc tgcctttgcg ccgcggccac ctctggagta ccgccattt ctgttgccgt 30000
 35
 cagatggcaa aaacactatc gccacottgt ttgccttctc cttgtaggcg tattttttct 30060
 caatgtctc tatcacctgg gaaatcagg tgtggatata cacatttttt tggatgtcca 30120
 40
 ggctaaagct tccgctttcg attctgtca cctgcagaat gctttcggca aggttctgca 30180
 gccgggaagc gtttcttgtt atcatgtcaa gctcccgctg aaactctgtt tttctttcgc 30240

caagcttctc ctccagtatc tccacaccgt ttaggatggg catgattggc gttcgcaact 30300

catgcgcagc cacgtttatg aactcgcttt tgactttgtc gttttgggtca agctgctgga 30360

5

acaggacact ctggtcatag aggacctcaa atatggacga gtaagacaat accgttggct 30420

cgctgttgga gtagattgaa aagccgattg cggcggttgc cacctcctcc cttgcgtgta 30480

10

tcagctccat caccagcgac tcctttctgt ccacaaccag tgtctttatc ttgatgcaa 30540

tgcttggcgc aatgtcctgg acttggatgt tgggcctgta ttttgtgagg agcctcaagg 30600

15

acaaggactc tcgcactgag gcatccatcg gcgtgaggat gttgatcctc aggctgtcgt 30660

tttgcctcac catotccttc aagagttgca gtgtgccgcc tttttcctgc aggtggaacg 30720

cgtaaccgt ggagtacatt atcagtatct cccttttggc cctgcttatc atttcaaact 30780

20

ccctttggac cgogtccttg tagttggaga agaccagtga gacgggcatg acaaccccat 30840

cotccagctc ctttatcctg tgctctgcgg gcaacgccct gcccctaaag ctgtcaaaca 30900

25

caaactgctg ctgctctgca atctcagga ggttctgtaa caaagctgg ggtattgact 30960

gtgccgcatg aagggtggtg acagccacat actccctctg gtcggccacc tcaaagtctc 31020

30

ccttcagccc atccaggtgc ctaatctccg aaaacgagag catctccttg acatagccga 31080

cgttgtcctt tgttatttcg gttacatacc gcagtttaag gccctggtt ttgaccgcgt 31140

caaccctttt ctcccttatg gcgtcaaccc ctatcatcac ggacggggcc aaggagtta 31200

35

tgcaagagtc tatcttcaca ttggccctgt ctatgaacct caaaatggcg ttgtttgcgt 31260

tttcagggcc atagtacacc tttgtggtgg gtgcaccggc gtcaacacca ccatcattat 31320

40

cgttctcgtt attgttgacg tcacaaaaac cgcgtttgtg cgacaaaccg gggtttgctg 31380

caatgtattt attttcacat acattatcat tatttgtgcc aatgtcatgg agattattat 31440

tatgtattag agaagaacct tccaaccgtt atcacgatat cctattttgt tcactatatt 31500
 aatttgagct taaaacttta taaataccgt atatacggta acaggattat tattctaaaa 31560
 5 aaacacttaa aggtacttga caaaattctg aacaaaagat ccccatatt tactaaatac 31620
 caagatttct gcaaatcgat gtgatgtgat gtatgcatag taccaatatc taggcaaacg 31680
 tttttggcat tagaaaggaa tacgaataga taatcaaaga atgaaatggc cgaacacaaa 31740
 10 ccaacaccat ggctttcaca cactgtcatt attatgtgaa actttatctg gcctctaatac 31800
 tttgtcagga attaaactgt tttttattgc caattctata atgatatgct ataagcagtt 31860
 15 agattacctt ttgatggtag tggttgttcc agtagtggtt tctccagtaa tatcatgagt 31920
 ttaaagaccc ggctgatggc agcgatagaa tgcttaattg catctattga ggaaagtgtt 31980
 gttgaaggta caacgctcaa tactatcaat tgaggacaac agggattgag aattgttttg 32040
 20 acaatgataa tccattcata aaaaaaattg caagataaag catatgccgc gattgttgac 32100
 cccctatttt gcatgcgttc caacaaaaag ttgtcttaac tttgcagaca tttgaataaa 32160
 25 ttaaaaagat gttgttgact ttcgtttatt gattgattaa gattacggtt ttattttacc 32220
 aaggatttaa gcattacttg cctacacgaa attaaattgc gagcaggaaa acaggaatgt 32280
 gtttacataa taagaatata cccctaacca agtctttttc ctatcgcatc tttttttggc 32340
 30 tacgccaggg cgaagaatat acttttggtt acaatgattg gtaaaccctt taaccttgct 32400
 tttgcgtgaa ttgtcataat tgatgttcgt aaagataaaa gcaataaaaa gaaatagtca 32460
 35 ttatgtagaa taacacattt tttttataac ccgttataat ttaattgcaa agcagtcatac 32520
 tttctaaaat aatcacaatt tgcagaatgc cgtcacttca tcttggttgca tatggtttaa 32580
 ttttgatata tttcgaaagc ccaatcacaa gggttaaacgg tagaacaagt cacttgatta 32640
 40 ttaaaatata tccacatatg gataacaata caaggatgag ttcttttagca atcgagtttt 32700

ttttatccct tttttcaata acgttacttt ctaaaagaat ataccaacca gtgaaatcaa 32760
 agtcatatac ctacatgac aagcatccat ttcagtacaa gatggaggat tatgcaaacc 32820

 5 acaacaaaat tgtagactat aaaaactgct tacttttttt tcaagtatcg atgttacaaa 32880
 aaaataaaat aattaggatt cgggttccag gtttgtttta tacaggtggc tggatttccc 32940
 tcacactaaa gtttttgata tccacatcat ttgcaccatc ccacctgaaa gtagcaatgg 33000
 10 ggctcccca ggatataatc tgatccggct caccaccaca ctcttcacca tcatttccaa 33060
 acccactga gtcagtgaat gtgtatacct tttgccaatt gttcttcaga gtcgggctat 33120
 15 ccgggtttct gtctacccat atttcagtgg tgactacggc ctcaccacca gccaatgggt 33180
 ggttatagat catggcttta aatccaatga atctatcaaa actagacgcc gaaggtgagg 33240
 gtgtggtagt gcttgaaaac acataggaga catgccactg ctcttttgca agcctaacc 33300
 20 ttccatcata gaatagatct gctttatatg ctgagccctc gcctccttcg ccatcatagt 33360
 gctaccacc cctgtcatac caagcgaaat tttcagaatc atctccacta ttaaccctta 33420
 25 caataccgct catttccaca ttcttccaat catttgata ctgcatgtat ccttgtgttg 33480
 cgagtaccga gtgatcgtaa gtctcaatat cctctggatg gtaccctgat gatgtaaaca 33540
 cgttatatct gacctgatcg tcattaacgt tccaactgcc atctgggttt aggtccatgt 33600
 30 caggtgggtt tgttcgtgga tcattgttcg ggttttgcat attcataaac catttttctc 33660
 caccaccgc cttatcgggg taaatctggg ttatcccaa ctggtcta atgtccctccgc 33720
 35 ctccagaagg ggcaacagag aacgtccata ccttgctggc agccaatggg acaccagtcg 33780
 catccgtagc accggttggt attctggcag tgtatgtggc accaggtgtt aaatctgcag 33840
 aggggttttag ggtcgcaact gtgttggttg gtgaattcat gcttacggtt gcgggcacag 33900
 40 gtgcgcctcc gcttggttagc agtgt 33925

<210> 2

<211> 2367

5

<212> DNA

<213> Crenarchaeote

10

<400> 2

	atggtgataa aatccgactt gcctctggag gagaaaaata gttattatca aaaagaactt	60
15	ccagaaaata ttccatcatt gttactttct tccgtttaca taggagaaaa aaaatcagtg	120
	tttttgaagt ttacaatcc agaagattct caaatatatt ttggagcga gtcttttatt	180
	gaaaatcata taaataaaca tcaaccttat tgctttgtaa aggaactcta ttctgatcag	240
20	gttaaaacaa tagttagcaa agagccacat aggttttagac tagaaaaaat aaaaaaaatg	300
	gacgatattg aggacaagga aatatcggtt tttaaaataa ttgccctga ccccttttcc	360
25	attggtggaa cagatagtag ttttagggaa aaggttactt cttgggaggc tgatatcaaa	420
	taccatgaaa gttattttatt cgatttgggt ctaattccgg gagcgtttta taacagaata	480
	ggcaataatt tagtttttca tgaattccca atgccagaga aagttgacga atatctggat	540
30	aatctcataa aacctaattt taaagaaaat gagccaaaaa gtagtgaata taatgagttt	600
	ctaataaagt ggtcaagatt gttaaaccag cctattccgg atatcaaaag aatttctttg	660
35	gatatagaag tggactctga agaggggaagg atgccacag ccagagatca cgataaagta	720
	attactgcag tgggtttatc ggcacoggat ggatttagaa aggtattcgt cttaagaaaa	780
40	gatccaaatt ttgatccctc taaactagat tcaacaaccg ttgaattatg tgatagcgaa	840
	aaagacatga tactaaaagt ttttgctatt attcaaaatt atccaatagt tttaaccttt	900

aatggtgatg attttgattt accttattta tatgctagat ctcaagaccc atcgatagac 960

cctgtacaca aaaaacccat tagtaaagaa ttggtgccta ttttagttaa aaaagactct 1020

5 tttataaaaa ggggtattca ggcggtaccg gtttccttaa agcatggaat ccatatogat 1080

ttattcagga catttcaaaa taaatctgta cagaattatg ctttttagtca taaatactct 1140

10 gagtttactt taaatgctat ctgcgaagcc ctattaaacg agtcaaaaat agactttgat 1200

gaaagcatag gtgatcttcc attggaaaaa ctggccgagt attgcctcaa agatgcagac 1260

15 ttgacatttc gtctgacatc tttcaatgac aatttactga taaaattggt gattatcatt 1320

tctaggatat cccgaatgtc aatagaagat ataacaagat tcggggtgaa tcaatggatt 1380

aggtccatga tgttttttga acataggcag caaaatatca ttattccccg taaagatgaa 1440

20 ttacagaaaa aaggaacatc gtctacagtt gccattataa aggaaaaaaa atatcgagga 1500

gggtctggtg ttgagcccg ttttaggaatt catttcaatg tcatagttgt agattttgct 1560

agtctgtatc ctagcataat taaagttcac aatttatctt acgaaacagt caattgtcct 1620

25 catgaaaatt gcagaaggga tccatcaaca catattgagc aaacaaacca ttgggtttgc 1680

aaggaaaagc aagggatgac ctccatattg ataggaaccc taagggatct aagggttaat 1740

30 tattacaaat atctatcaaa ggataattct ttggataaag aggataaaca gctatacagt 1800

gttatcagtc aggccataaa ggttatttta aatgctacgt atggggttat ggggtgctgaa 1860

atatttccgc tctattgttt acctgtagct gaggctaccg cagcggttg aaggatgacc 1920

35 acaacaaaaa ctattgaaaa atgcaacgaa gaaaagattg aggttattta cggtgatacg 1980

gattctctgt tcctaaagaa tccttccaag gaaggattaa gtggaatttc atcctggtct 2040

40 aaaaaagaac taggcataga tttggagata gataaaagat atcgctacgt ggtttttagt 2100

gaactaaaaa aaaattacct aggtgtattg gaggacggaa ctgtagatgt taaaggatta 2160

acaggggaaga agtctcatatc acctccaata ataagacaag ctttctatga catattaaat 2220

gtccttaag aaatatcttc agaaaaagac tttgaaagag caaaggaaaa gataaaaaaa 2280

5

atagtgaat caattgcaga aaacttgag aaaaaagaa tttctctgga agaattaagt 2340

tttaatgtta tgatcaacaa ggctgtg 2367

10

<210> 3

<211> 789

15

<212> PRT

<213> Crenarchaeote

20

<400> 3

Met	Val	Ile	Lys	Ser	Asp	Leu	Pro	Leu	Glu	Glu	Lys	Asn	Ser	Tyr	Tyr
1				5					10					15	

25

Gln	Lys	Glu	Leu	Pro	Glu	Asn	Ile	Pro	Ser	Leu	Leu	Leu	Ser	Ser	Val
			20					25					30		

30

Tyr	Ile	Gly	Glu	Lys	Lys	Ser	Val	Phe	Leu	Lys	Phe	Tyr	Asn	Pro	Glu
		35					40					45			

35

Asp	Ser	Gln	Ile	Tyr	Phe	Trp	Ser	Glu	Ser	Phe	Ile	Glu	Asn	His	Ile
		50				55					60				

40

Asn	Lys	His	Gln	Pro	Tyr	Cys	Phe	Val	Lys	Glu	Leu	Tyr	Ser	Asp	Gln
65					70					75				80	

Val Lys Thr Ile Val Ser Lys Glu Pro His Arg Phe Arg Leu Glu Lys
 85 90 95

5 Ile Lys Lys Met Asp Asp Ile Glu Asp Lys Glu Ile Ser Val Phe Lys
 100 105 110

10 Ile Ile Ala Pro Asp Pro Leu Ser Ile Gly Gly Thr Asp Ser Ser Phe
 115 120 125

15 Arg Glu Lys Val Thr Ser Trp Glu Ala Asp Ile Lys Tyr His Glu Ser
 130 135 140

Tyr Leu Phe Asp Leu Gly Leu Ile Pro Gly Ala Phe Tyr Asn Arg Ile
 145 150 155 160

20 Gly Asn Asn Leu Val Phe His Glu Phe Pro Met Pro Glu Lys Val Asp
 165 170 175

25 Glu Tyr Leu Asp Asn Leu Ile Lys Pro Asn Phe Lys Glu Asn Glu Pro
 180 185 190

30 Lys Ser Ser Glu Tyr Asn Glu Phe Leu Ile Lys Trp Ser Arg Leu Leu
 195 200 205

35 Asn Gln Pro Ile Pro Asp Ile Lys Arg Ile Ser Leu Asp Ile Glu Val
 210 215 220

Asp Ser Glu Glu Gly Arg Met Pro Thr Ala Arg Asp His Asp Lys Val
 225 230 235 240

40 Ile Thr Ala Val Gly Leu Ser Ala Ser Asp Gly Phe Arg Lys Val Phe
 245 250 255

Val Leu Arg Lys Asp Pro Asn Phe Asp Pro Ser Lys Leu Asp Ser Thr
 260 265 270

5

Thr Val Glu Leu Cys Asp Ser Glu Lys Asp Met Ile Leu Lys Val Phe
 275 280 285

10

Ala Ile Ile Gln Asn Tyr Pro Ile Val Leu Thr Phe Asn Gly Asp Asp
 290 295 300

15

Phe Asp Leu Pro Tyr Leu Tyr Ala Arg Ser Gln Asp Pro Ser Ile Asp
 305 310 315 320

20

Pro Val His Lys Lys Pro Ile Ser Lys Glu Leu Val Pro Ile Leu Val
 325 330 335

25

Lys Lys Asp Ser Phe Ile Lys Arg Gly Ile Gln Ala Asp Pro Val Ser
 340 345 350

30

Leu Lys His Gly Ile His Ile Asp Leu Phe Arg Thr Phe Gln Asn Lys
 355 360 365

35

Ser Val Gln Asn Tyr Ala Phe Ser His Lys Tyr Ser Glu Phe Thr Leu
 370 375 380

Asn Ala Ile Cys Glu Ala Leu Leu Asn Glu Ser Lys Ile Asp Phe Asp
 385 390 395 400

40

Glu Ser Ile Gly Asp Leu Pro Leu Glu Lys Leu Ala Glu Tyr Cys Leu
 405 410 415

Lys Asp Ala Asp Leu Thr Phe Arg Leu Thr Ser Phe Asn Asp Asn Leu
 420 425 430

5 Leu Ile Lys Leu Leu Ile Ile Ile Ser Arg Ile Ser Arg Met Ser Ile
 435 440 445

10 Glu Asp Ile Thr Arg Phe Gly Val Asn Gln Trp Ile Arg Ser Met Met
 450 455 460

15 Phe Phe Glu His Arg Gln Gln Asn Ile Ile Ile Pro Arg Lys Asp Glu
 465 470 475 480

Leu Gln Lys Lys Gly Thr Ser Ser Thr Val Ala Ile Ile Lys Glu Lys
 485 490 495

20 Lys Tyr Arg Gly Gly Leu Val Val Glu Pro Val Leu Gly Ile His Phe
 500 505 510

25 Asn Val Ile Val Val Asp Phe Ala Ser Leu Tyr Pro Ser Ile Ile Lys
 515 520 525

30 Val His Asn Leu Ser Tyr Glu Thr Val Asn Cys Pro His Glu Asn Cys
 530 535 540

35 Arg Arg Asp Pro Ser Thr His Ile Glu Gln Thr Asn His Trp Val Cys
 545 550 555 560

Lys Glu Lys Gln Gly Met Thr Ser Ile Leu Ile Gly Thr Leu Arg Asp
 565 570 575

40 Leu Arg Val Asn Tyr Tyr Lys Tyr Leu Ser Lys Asp Asn Ser Leu Asp
 580 585 590

Lys Glu Asp Lys Gln Leu Tyr Ser Val Ile Ser Gln Ala Ile Lys Val
 595 600 605

5

Ile Leu Asn Ala Thr Tyr Gly Val Met Gly Ala Glu Ile Phe Pro Leu
 610 615 620

10

Tyr Cys Leu Pro Val Ala Glu Ala Thr Ala Ala Val Gly Arg Met Thr
 625 630 635 640

15

Thr Thr Lys Thr Ile Glu Lys Cys Asn Glu Glu Lys Ile Glu Val Ile
 645 650 655

20

Tyr Gly Asp Thr Asp Ser Leu Phe Leu Lys Asn Pro Ser Lys Glu Gly
 660 665 670

25

Leu Ser Gly Ile Ser Ser Trp Ser Lys Lys Glu Leu Gly Ile Asp Leu
 675 680 685

30

Glu Ile Asp Lys Arg Tyr Arg Tyr Val Val Phe Ser Glu Leu Lys Lys
 690 695 700

35

Asn Tyr Leu Gly Val Leu Glu Asp Gly Thr Val Asp Val Lys Gly Leu
 705 710 715 720

Thr Gly Lys Lys Ser His Thr Pro Pro Ile Ile Arg Gln Ala Phe Tyr
 725 730 735

40

Asp Ile Leu Asn Val Leu Lys Glu Ile Phe Ser Glu Lys Asp Phe Glu
 740 745 750

Arg Ala Lys Glu Lys Ile Lys Lys Ile Val Gln Ser Ile Ala Glu Asn
 755 760 765

5 Leu Glu Lys Lys Arg Ile Ser Leu Glu Glu Leu Ser Phe Asn Val Met
 770 775 780

10 Ile Asn Lys Ala Val
 785

<210> 4

15 <211> 882

<212> DNA

20 <213> Crenarchaeote

<400> 4

25 atgggatattg atcataaaat tttagatatat ttcatattat ctattaacaa aataattatt 60
 acaatggggtt tggtttcgga tagacaaaga aacgagacaa tggattttat aaaaatactg 120
 ggatataaca tcagatatat aaaaatagat caagtcaagt caaatgaaac cataattctg 180
 30 cttcatggta taggagcttc cgcagaacga tggtcagaat tagtcccatt tttgtataat 240
 tgcaatataa ttataccaga catcattggt tttgggttaca gtgaaaaacc aaggatagag 300
 35 tacaacatag atttatttgt aaagtttttg gatgaattgt ttctgaaact tgaaatcaaa 360
 aaccccataa taatgggttc gtottttgggt ggtcaattga ttttagaata ttatttcagg 420
 cacaaagact tttttaaaaa aatgattcta gtgtccccgg ccggtaccca agagagaccg 480
 40 aactagcgt taaggcaata cacttactca tgtttatacc caacaagaga aaataccgaa 540

agagcattta agatgatgtc gcatttcaat cacacagtaa aagattcaat gataaaggat 600

tttattaata gaatgaagca gcccaacgca aaacactcgt ttgtttcaac acttttagca 660

5 ctaaggaaaa atagtgattt acaagacaac ctgagggaaa tcaaaatccc aactttagta 720

atatggggaa aagaggacaa caccattcca gtagaaaata tagagtattt caggggcatc 780

ccttttgtaa aaacatgcat aatgagtgat tgcggtcatg tgccttttgt tgaaaagcct 840

10

cttgagtttt ataaaatagt caaagagttt atcgactcct aa 882

<210> 5

15

<211> 293

<212> PRT

20

<213> Crenarchaeote

<400> 5

25

Met	Asp	Ile	Asp	His	Lys	Ile	Leu	Val	Tyr	Phe	Ile	Leu	Ser	Ile	Asn
1				5					10					15	

30

Lys	Ile	Ile	Ile	Thr	Met	Gly	Leu	Val	Ser	Asp	Arg	Gln	Arg	Asn	Glu
			20					25						30	

35

Thr	Met	Asp	Phe	Ile	Lys	Ile	Leu	Gly	Tyr	Asn	Ile	Arg	Tyr	Ile	Lys
		35						40					45		

40

Ile	Asp	Gln	Val	Lys	Ser	Asn	Glu	Thr	Ile	Ile	Leu	Leu	His	Gly	Ile
	50					55						60			

Gly	Ala	Ser	Ala	Glu	Arg	Trp	Ser	Glu	Leu	Val	Pro	Phe	Leu	Tyr	Asn
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

65

70

75

80

Cys Asn Ile Ile Ile Pro Asp Ile Ile Gly Phe Gly Tyr Ser Glu Lys

5

85

90

95

Pro Arg Ile Glu Tyr Asn Ile Asp Leu Phe Val Lys Phe Leu Asp Glu

100

105

110

10

Leu Phe Leu Lys Leu Glu Ile Lys Asn Pro Ile Ile Met Gly Ser Ser

115

120

125

15

Phe Gly Gly Gln Leu Ile Leu Glu Tyr Tyr Phe Arg His Lys Asp Phe

130

135

140

20

Phe Lys Lys Met Ile Leu Val Ser Pro Ala Gly Thr Gln Glu Arg Pro

145

150

155

160

Thr Leu Ala Leu Arg Gln Tyr Thr Tyr Ser Cys Leu Tyr Pro Thr Arg

25

165

170

175

Glu Asn Thr Glu Arg Ala Phe Lys Met Met Ser His Phe Asn His Thr

180

185

190

30

Val Lys Asp Ser Met Ile Lys Asp Phe Ile Asn Arg Met Lys Gln Pro

195

200

205

35

Asn Ala Lys His Ser Phe Val Ser Thr Leu Leu Ala Leu Arg Lys Asn

210

215

220

40

Ser Asp Leu Gln Asp Asn Leu Arg Glu Ile Lys Ile Pro Thr Leu Val

225

230

235

240

Ile Trp Gly Lys Glu Asp Asn Thr Ile Pro Val Glu Asn Ile Glu Tyr
 245 250 255

5

Phe Arg Gly Ile Pro Phe Val Lys Thr Cys Ile Met Ser Asp Cys Gly
 260 265 270

10

His Val Pro Phe Val Glu Lys Pro Leu Glu Phe Tyr Lys Ile Val Lys
 275 280 285

15

Glu Phe Ile Asp Ser
 290

<210> 6

20

<211> 318

<212> DNA

<213> Crenarchaeote

25

<400> 6

ttgaatcaat ccacttctat gagtaatgag aatgaagaaa ataaagatat agattttaag 60

30

aaatccattg aaaaggctgc ggaattccag caggatttgt tgcgacagtt ctctacaatt 120

caatacaatg cgtttcagaa tatgttttca tctttgcaag gatttacaaa ttataatgcc 180

35

atgttttaaaa ccaccgtaca gacgggtggc aggatctcaa ttcccgaagc agaaagaaat 240

gctttgggga ttgaagaggg tgatctagtc caggttataa ttataccggt gacaaggaaa 300

aagaaaaaca caagttaa 318

40

<210> 7

<211> 105

5 <212> PRT

<213> Crenarchaeote

10

<400> 7

Met Asn Gln Ser Thr Ser Met Ser Asn Glu Asn Glu Glu Asn Lys Asp
 1 5 10 15

15

Ile Asp Phe Lys Lys Ser Ile Glu Lys Ala Ala Glu Phe Gln Gln Asp
 20 25 30

20

Leu Leu Arg Gln Phe Ser Thr Ile Gln Tyr Asn Ala Phe Gln Asn Met
 35 40 45

25

Phe Ser Ser Leu Gln Gly Phe Thr Asn Tyr Asn Ala Met Phe Lys Thr
 50 55 60

30

Thr Val Gln Thr Gly Gly Arg Ile Ser Ile Pro Glu Ala Glu Arg Asn
 65 70 75 80

35

Ala Leu Gly Ile Glu Glu Gly Asp Leu Val Gln Val Ile Ile Ile Pro
 85 90 95

Leu Thr Arg Lys Lys Lys Asn Thr Ser
 100 105

40

<210> 8

<211> 1086

<212> DNA

5 <213> Crenarchaeote

<400> 8

10 atgagaaaaa aaatgaataa ttccttaatt aattatttag tgaatgatta ttttacgttt 60

gttagggatc ctgataacat ttcaaaatta aaagaactca ggaagaaatt gtcgaatata 120

15 gaaaacataa agactggatc aagcgaatat gaggtaataa gggaaacaac cctcttcggt 180

ttactacatt ataaaccctt aaagcaacaa actttcaagt accctttggt gattgtttat 240

gcattaataa acaaatcata tattttggat ctgcagaacg acaaaagttg gataaggaac 300

20 ctgctagagc agggcataaa tgtctatctg attgactgga aacccccgtc aaaactggat 360

aaatacatca ctgttgatga ttatgtcaat ttgtttattt atgagtgtgt agaatacata 420

25 aaaaacatag aaaacattga tcagatttca ttacaaggat attgcatggg ggttacaatg 480

tccttgatgt acaattcgct atatcaaaaa aacattaata atctagtcac cattgctcca 540

attgttgatg ccgagaaaga caaatccgta ataaaaaaca tggctgagca catggatatt 600

30 gacaaagtac tgtcctatca cgaaaacttt ccatatgaat tactgtatct ggtttatgca 660

tcactaaaac cattcaagca aggtgtaaac aaatactata atttatttaa aaactttgaa 720

gatgaaagtt ttgtacagaa ctttttaaga atagagaaat ggctgtatga cacacctcct 780

35 attgcggggg aaacctttag gcaatgggta aaggatatct atcagcaaaa cctttttgca 840

aaaaacaaga tgattgtggg tgaaaacaag ataaatttgt caaacattaa ggttcccgtt 900

40 cttaatgttg tagctgaatt tgaccacctt gtaacgtctg acagcagtag ctccctaaac 960

aacctaattt caagtcagga taaaagcctg atgaaatttc caacagggca tgtagggcta 1020

attgctagca acttttcaca gaaaaatgtt ttaccaaaaa ttggaaaatg gattcaaaca 1080

cattaa

1086

5

<210> 9

<211> 361

10

<212> PRT

<213> Crenarchaeote

15

<400> 9

Met Arg Lys Lys Met Asn Asn Ser Leu Ile Asn Tyr Leu Val Asn Asp
20 1 5 10 15

Tyr Phe Thr Phe Val Arg Asp Pro Asp Asn Ile Ser Lys Leu Lys Glu
25 20 25 30

Ile Arg Lys Lys Leu Ser Asn Ile Glu Asn Ile Lys Thr Gly Ser Ser
35 40 45

30

Glu Tyr Glu Val Ile Arg Glu Thr Thr Leu Phe Arg Leu Leu His Tyr
50 55 60

Lys Pro Leu Lys Gln Gln Thr Phe Lys Tyr Pro Leu Leu Ile Val Tyr
35 65 70 75 80

Ala Leu Ile Asn Lys Ser Tyr Ile Leu Asp Leu Gln Asn Asp Lys Ser
40 85 90 95

Trp Ile Arg Asn Leu Leu Glu Gln Gly Ile Asn Val Tyr Leu Ile Asp
 100 105 110

5 Trp Lys Pro Pro Ser Lys Leu Asp Lys Tyr Ile Thr Val Asp Asp Tyr
 115 120 125

10 Val Asn Leu Phe Ile Tyr Glu Cys Val Glu Tyr Ile Lys Asn Ile Glu
 130 135 140

15 Asn Ile Asp Gln Ile Ser Leu Gln Gly Tyr Cys Met Gly Gly Thr Met
 145 150 155 160

Ser Leu Met Tyr Thr Ser Leu Tyr Gln Lys Asn Ile Lys Asn Leu Val
 165 170 175

20 Thr Ile Ala Pro Ile Val Asp Ala Glu Lys Asp Lys Ser Val Ile Lys
 180 185 190

25 Asn Met Ala Glu His Met Asp Ile Asp Lys Val Leu Ser Tyr His Glu
 195 200 205

30 Asn Phe Pro Tyr Glu Leu Leu Tyr Leu Val Tyr Ala Ser Leu Lys Pro
 210 215 220

35 Phe Lys Gln Gly Val Asn Lys Tyr Tyr Asn Leu Phe Lys Asn Phe Glu
 225 230 235 240

Asp Glu Ser Phe Val Gln Asn Phe Leu Arg Ile Glu Lys Trp Leu Tyr
 245 250 255

40 Asp Thr Pro Pro Ile Ala Gly Glu Thr Phe Arg Gln Trp Val Lys Asp
 260 265 270

Ile Tyr Gln Gln Asn Leu Phe Ala Lys Asn Lys Met Ile Val Gly Glu
 275 280 285

5

Asn Lys Ile Asn Leu Ser Asn Ile Lys Val Pro Val Leu Asn Val Val
 290 295 300

10

Ala Glu Phe Asp His Leu Val Thr Ser Asp Ser Ser Ser Ser Leu Asn
 305 310 315 320

15

Asn Leu Ile Ser Ser Gln Asp Lys Ser Leu Met Lys Phe Pro Thr Gly
 325 330 335

20

His Val Gly Leu Ile Ala Ser Asn Phe Ser Gln Lys Asn Val Leu Pro
 340 345 350

Lys Ile Gly Lys Trp Ile Gln Thr His
 355 360

25

<210> 10

<211> 582

30

<212> DNA

<213> Crenarchaeote

35

<400> 10

ttgcaattag aaaataacaa tattggagag gaaaaaaaca gtaaaaacac tctatctgaa 60

40

gaggcaggac ttcagtctgt atttgaaaac ttataaaaac aattaacaga gttaaatagc 120

cttacaacct tggggccatt cacctcttta atgaatgatc caaaccttaa tttaaataca 180

ttaaaggaac acggtaatTT gttactgaga tatcagtcac ttctcaacct atacttttcc 240
 cgtatgataa atgcttattt gttggccgta aacaaggat -cgtctgctat agatgaaaaa 300
 5 aaccccgacg atattaggaa aataatcata aatacttttg aggatgtgtt ctcgtcaatg 360
 ttgcagtcaa cagacttttc aatcaattat aacaatttat tgaattccag cattgatgtc 420
 10 atcaaaagtt atcaaaaaat ttacgattca aatgccgttt tgttttaggtc acaacaacaa 480
 ctgtcaaaag aagaaaaaga cctggtattt tataatctct atgaaatcaa aaaaatatca 540
 ttggaaatca aaaaaaaatt aaatgagaaa aaaaatgaat aa 582

15

<210> 11

<211> 193

20

<212> PRT

<213> Crenarchaeote

25

<400> 11

30

Met Gln Leu Glu Asn Asn Asn Ile Gly Glu Glu Lys Asn Ser Lys Asn
 1 5 10 15

Thr Leu Ser Glu Glu Ala Gly Leu Gln Ser Val Phe Glu Asn Phe Ile
 20 25 30

35

Lys Gln Leu Thr Glu Leu Asn Ser Leu Thr Thr Leu Gly Pro Phe Thr
 35 40 45

40

Ser Leu Met Asn Asp Pro Asn Leu Asn Leu Asn Thr Leu Lys Glu His
 50 55 60

Gly Asn Leu Leu Leu Arg Tyr Gln Ser Phe Leu Asn Leu Tyr Phe Ser
 65 70 75 80

5

Arg Met Ile Asn Ala Tyr Leu Leu Ala Val Asn Lys Val Ser Ser Ala
 85 90 95

10

Ile Asp Glu Lys Asn Pro Asp Asp Ile Arg Lys Ile Ile Ile Asn Thr
 100 105 110

15

Phe Glu Asp Val Phe Ser Ser Met Leu Gln Ser Thr Asp Phe Ser Ile
 115 120 125

20

Asn Tyr Asn Asn Leu Leu Asn Ser Ser Ile Asp Val Ile Lys Ser Tyr
 130 135 140

25

Gln Lys Ile Tyr Asp Ser Asn Ala Val Leu Phe Arg Ser Gln Gln Gln
 145 150 155 160

30

Leu Ser Lys Glu Glu Lys Asp Leu Leu Phe Tyr Asn Leu Tyr Glu Ile
 165 170 175

35

Lys Lys Ile Ser Leu Glu Ile Lys Lys Lys Leu Asn Glu Lys Lys Asn
 180 185 190

Glu

40 <210> 12

<211> 438

<212> DNA

5 <213> Crenarchaeote

<400> 12

10 atgcctacaa gttcagatgt tttatacatg tccaaaccag cgggtggtatg tatacattct 60
 tttgacatgg tgggcgggta tgcgcatacc caaaaactaa gatgctgtat cagcctcggg 120
 aagaggggta tgtggggggac aatagaaaga atccatccac aaacgaatgg ttttggcaaa 180
 15 tgtctgcctt ggctcattgt ttctatatat ggtttcgcca tagataatat ttgggtaatt 240
 actcatgtgc ctgtatacaa aaatagacaa ccatctctac ctatatataa attttttgac 300
 20 aagggtcagg ttcttttttct cttttttttg caaattatgc cgggccaccc agaaacaaac 360
 cccgcctttg ggcccgcaat gattactggg caacccaaat ctagcgcccc acgcccagga 420
 gtgcaagtga caatgtga 438

25

<210> 13

<211> 145

30

<212> PRT

<213> Crenarchaeote

35

<400> 13

Met Pro Thr Ser Ser Asp Val Leu Tyr Met Ser Lys Pro Ala Val Val

40

1

5

10

15

Cys Ile His Ser Phe Asp Met Val Gly Gly Tyr Ala His Thr Gln Lys
 20 25 30

5 Leu Arg Cys Cys Ile Ser Leu Gly Lys Arg Val Met Trp Gly Thr Ile
 35 40 45

10 Glu Arg Ile His Pro Gln Thr Asn Gly Phe Gly Lys Cys Leu Pro Trp
 50 55 60

15 Leu Ile Val Ser Ile Tyr Gly Phe Ala Ile Asp Asn Ile Trp Val Ile
 65 70 75 80

Thr His Val Pro Val Tyr Lys Asn Arg Gln Pro Ser Leu Pro Ile Tyr
 85 90 95

20 Lys Phe Phe Asp Lys Gly Gln Val Leu Phe Leu Leu Phe Leu Gln Ile
 100 105 110

25 Met Pro Gly His Pro Glu Thr Asn Pro Ala Phe Gly Pro Ala Met Ile
 115 120 125

30 Thr Gly Gln Pro Lys Ser Ser Ala Pro Arg Pro Gly Val Gln Val Thr
 130 135 140

Met
 145

35

<210> 14

<211> 915

40

<212> DNA

<213> Crenarchaeote

5 <400> 14
 ttggattctt ggggcgaatc taatattgtc ctttggctgc tacttaggct attcaagcca 60
 aaaaccaaga tctttgttgt gtttcatcac catgaaccgc ggatatctat ttgcaagaac 120
 10 tattttgaat ttctgtacaa ttatctaata caaaaggcta ctgcggtgat gcttaaggat 180
 tctgatatga ttttgaccgt gagtcaagcg tcaaagcatg aactcaacac agtctatgga 240
 ataggggtta gcaaaatcaa taatttgaag gaaacagcaa ataaaaaac cagggaatta 300
 15 gcaaaaaatc tgaccaacag aattgccatt gtaggaactg gaatagataa aaatatcttt 360
 ttaaaggatt ccaacagagg agtaatcaac aataaaaagg acattgattt tctttgtatc 420
 20 ggaaggatag aaaaatttca tggactggag gaaatttga ctgcaataaa aacactcaga 480
 ccagaatcta attttgtaat ggttgggcgc ataccacctg ataaggctgc aaaactacgt 540
 aatgcgggta tagatcacag aggctttgtc tccgaggaag aaaagattag cctttattct 600
 25 aaatctaaag tctttatttt tccatcatcc agagagggtt ttggcattgc tgtggctgag 660
 gccttagttt cgtgtgttcc cactgttgcc tggaaaactcc ccgtttttga agaactatac 720
 30 ttaaaaaatg gtaatacaaa cataaaacta atagaatatg gagaaaccac cctgtttgca 780
 gaagagtgcg taaaaatgct aaataaatat ggcataatca aaaaggcgac tgaaggaaaa 840
 aaggtcagtt tccaactccc aaactggcag acagtggcaa aaaatgtaat gacaacaata 900
 35 gaatctgtaa cctaa 915

<210> 15

40

<211> 304

<212> PRT

<213> Crenarchaeote

5

<400> 15

Met Asp Ser Trp Gly Glu Ser Asn Ile Val Leu Trp Leu Leu Leu Arg
 10 1 5 10 15

Leu Phe Lys Pro Lys Thr Lys Ile Phe Val Val Phe His His His Glu
 15 20 25 30

Pro Arg Ile Ser Ile Cys Lys Asn Tyr Phe Glu Phe Leu Tyr Asn Tyr
 35 40 45

20 Leu Ile Gln Lys Ala Thr Ala Val Met Leu Lys Asp Ser Asp Met Ile
 50 55 60

25 Leu Thr Val Ser Gln Ala Ser Lys His Glu Leu Asn Thr Val Tyr Gly
 65 70 75 80

30 Ile Gly Val Ser Lys Ile Asn Asn Leu Lys Glu Thr Ala Asn Lys Lys
 85 90 95

Thr Arg Glu Leu Ala Lys Asn Leu Thr Asn Arg Ile Ala Ile Val Gly
 100 105 110

35 Thr Gly Ile Asp Lys Asn Ile Phe Leu Lys Asp Ser Asn Arg Gly Val
 115 120 125

40 Ile Asn Asn Lys Lys Asp Ile Asp Phe Leu Cys Ile Gly Arg Ile Glu
 130 135 140

Lys Phe His Gly Leu Glu Glu Ile Trp Thr Ala Ile Lys Thr Leu Arg
 145 150 155 160

5

Pro Glu Ser Asn Phe Val Met Val Gly Arg Ile Pro Pro Asp Lys Ala
 165 170 175

10

Ala Lys Leu Arg Asn Ala Gly Ile Asp His Arg Gly Phe Val Ser Glu
 180 185 190

15

Glu Glu Lys Ile Ser Leu Tyr Ser Lys Ser Lys Val Phe Ile Phe Pro
 195 200 205

20

Ser Ser Arg Glu Gly Phe Gly Ile Ala Val Ala Glu Ala Leu Val Ser
 210 215 220

25

Cys Val Pro Thr Val Ala Trp Lys Leu Pro Val Phe Glu Glu Leu Tyr
 225 230 235 240

30

Leu Lys Asn Gly Asn Thr Asn Ile Lys Leu Ile Glu Tyr Gly Glu Thr
 245 250 255

35

Thr Leu Phe Ala Glu Glu Cys Val Lys Met Leu Asn Lys Tyr Gly Ile
 260 265 270

Ile Lys Lys Ala Thr Glu Gly Lys Lys Val Ser Phe Gln Leu Pro Asn
 275 280 285

40

Trp Gln Thr Val Ala Lys Asn Val Met Thr Thr Ile Glu Ser Val Thr
 290 295 300

<210> 16

<211> 1692

5

<212> DNA

<213> Crenarchaeote

10

<400> 16

atgtgtggaa ttgttggaat tttaagtaaa aaagagagaa atgttgcccc cttgatagga 60

15

aaaatgctat cctgtatgaa aaaccggggt cgggatggca tgggtttgtc tacagagaat 120

caaatagttt attctgatac ctttgataat ccattgtttt cacaggtaga ggggcatgac 180

20

gttttaggtc acagtcgttt ggcaatagtt ggtggctcct gtggtcagca gccgtttgtg 240

agttgtgata aaaaactcat tctggagcat aatggtgaaa tatataacta taaagaaatc 300

agaaagaacc tttctgcaca tcacactttt actacctcga ctgatagtga agttattgtt 360

25

caccttcttg aagaccatta tcaaaacact aaaggcgatc taatcgaagc tataaggaga 420

accgttacc agcttgatgg aatttatgtt ttggcgatta gagagcagtc cacaggagat 480

30

attgtgctgg tacgggatgg cattggagta agacaaattt actatgggtga aagtagtgat 540

ttcattgcat ttgcatcaga aagaaaagcc ttatggaaaa ttgctatgtc cgaccaaadc 600

aaaagacttt tgccaggcta tgctcttggtc atttcgogga aggaagggtc ctccaatttc 660

35

aagactacat tgtttcgat ttctgtaaat acaaaaaaat caatatgtga gaaatattca 720

atcctgtaca cagacatcga ttctgogggt aacgcatatg gtgatacatt ggttgaatct 780

40

atgagaaaac gtgtgagtga ctttaaaaaa atcgggtattg ttttctccgg tgggattgac 840

agtgtaatg tagcgtattt ggcaaaacaa atggcccccg aagttatttg ctatacgtct 900

gggattaaag gttcaagtga taccctcaac tcacttgaga tagcagaaaa acttgacctc . 960
 aagttggaaa tagaacagat gactgaaagt gatgttgaaa gtaccattcc aaaaataatc 1020
 5 agcataattg aagatgacaa catgggacag gttgaggttg ccattccaat atatggcgcg 1080
 gttaaattgg ctacgaaca ggaatacgg gtaatgctta caggtcaggg ggcagacgaa 1140
 ctgtttggcg gatattcctg gtattccaaa attgttaaaa aacacggata cgaaaaaatt 1200
 10 cagggatacc tgatagagga cattaagtta ctttacaag aaacactgga aagagaggac 1260
 aaaataacca tgtctcaaag catagagtta cgcaaccct ttttagatac taatctgata 1320
 15 gacacggtac tgagaataga tccgcgactc aatattcaaa acaatggcaa taactatgac 1380
 aacctaggaa aaagggttca ccgcaaactt gcagaaaaac tagggattcc aaaagagata 1440
 gcgtatagaa taaaggaagc agctcagcat gttctggga tacacaacac cctcaatact 1500
 20 ttggccatga aaaatgggtt tacggaatcc aagggttaatt ctagttatct ggacaaattg 1560
 aaaaaaaggg agcttatcgg cagctcacia agatatgggc atctttttga aaaggaacia 1620
 25 atctggagtt tggagccgca tatacagatg tatttggaga atatttcaaa aaacatattg 1680
 ccaaggaact ga 1692
 30
 <210> 17
 <211> 563
 35 <212> PRT
 <213> Crenarchaeote
 40
 <400> 17

Met Cys Gly Ile Val Gly Ile Leu Ser Lys Lys Glu Arg Asn Val Ala
 1 5 10 15

5 Pro Leu Ile Gly Lys Met Leu Ser Cys Met Lys Asn Arg Gly Pro Asp
 20 25 30

10 Gly Met Gly Leu Ser Thr Glu Asn Gln Ile Val Tyr Ser Asp Thr Phe
 35 40 45

15 Asp Asn Pro Leu Phe Ser Gln Val Glu Gly His Asp Val Leu Gly His
 50 55 60

Ser Arg Leu Ala Ile Val Gly Gly Ser Cys Gly Gln Gln Pro Phe Val
 65 70 75 80

20 Ser Cys Asp Lys Lys Leu Ile Leu Glu His Asn Gly Glu Ile Tyr Asn
 85 90 95

25 Tyr Lys Glu Ile Arg Lys Asn Leu Ser Ala His His Thr Phe Thr Thr
 100 105 110

30 Ser Thr Asp Ser Glu Val Ile Val His Leu Leu Glu Asp His Tyr Gln
 115 120 125

35 Asn Thr Lys Gly Asp Leu Ile Glu Ala Ile Arg Arg Thr Val Thr Gln
 130 135 140

Leu Asp Gly Ile Tyr Val Leu Ala Ile Arg Glu Gln Ser Thr Gly Asp
 145 150 155 160

40 Ile Val Leu Val Arg Asp Gly Ile Gly Val Arg Gln Ile Tyr Tyr Gly
 165 170 175

Glu Ser Ser Asp Phe Ile Ala Phe Ala Ser Glu Arg Lys Ala Leu Trp

180

185

190

5

Lys Ile Ala Met Ser Asp Gln Ile Lys Arg Leu Leu Pro Gly Tyr Ala

195

200

205

10

Leu Val Ile Ser Arg Lys Glu Gly Ser Ser Asn Phe Lys Thr Thr Leu.

210

215

220

15

Phe Pro Ile Ser Val Asn Thr Lys Lys Ser Ile Cys Glu Lys Tyr Ser

225

230

235

240

20

Ile Leu Tyr Thr Asp Ile Asp Ser Ala Val Asn Ala Tyr Gly Asp Thr

245

250

255

Leu Val Glu Ser Met Arg Lys Arg Val Ser Asp Phe Lys Lys Ile Gly

260

265

270

25

Ile Val Phe Ser Gly Gly Ile Asp Ser Val Ile Val Ala Tyr Leu Ala

275

280

285

30

Lys Gln Met Ala Pro Glu Val Ile Cys Tyr Thr Ser Gly Ile Lys Gly

290

295

300

35

Ser Ser Asp Ile Leu Asn Ser Leu Glu Ile Ala Glu Lys Leu Asp Leu

305

310

315

320

40

Lys Leu Glu Ile Glu Gln Met Thr Glu Ser Asp Val Glu Ser Thr Ile

325

330

335

Pro Lys Ile Ile Ser Ile Ile Glu Asp Asp Asn Met Gly Gln Val Glu
 340 345 350

5 Val Ala Ile Pro Ile Tyr Gly Ala Val Lys Leu Ala His Glu Gln Gly
 355 360 365

10 Ile Arg Val Met Leu Thr Gly Gln Gly Ala Asp Glu Leu Phe Gly Gly
 370 375 380

15 Tyr Ser Trp Tyr Ser Lys Ile Val Lys Lys His Gly Tyr Glu Lys Ile
 385 390 395 400

Gln Gly Tyr Leu Ile Glu Asp Ile Lys Leu Leu Tyr Lys Glu Thr Leu
 405 410 415

20 Glu Arg Glu Asp Lys Ile Thr Met Ser Gln Ser Ile Glu Leu Arg Glu
 420 425 430

25 Pro Phe Leu Asp Thr Asn Leu Ile Asp Thr Val Leu Arg Ile Asp Pro
 435 440 445

30 Arg Leu Asn Ile Gln Asn Asn Gly Asn Asn Tyr Asp Asn Leu Gly Lys
 450 455 460

35 Arg Val His Arg Lys Leu Ala Glu Lys Leu Gly Ile Pro Lys Glu Ile
 465 470 475 480

Ala Tyr Arg Ile Lys Glu Ala Ala Gln His Gly Ser Gly Ile His Asn
 485 490 495

40 Thr Leu Asn Thr Leu Ala Met Lys Asn Gly Phe Thr Glu Ser Lys Val
 500 505 510

Asn Ser Ser Tyr Leu Asp Lys Leu Lys Lys Arg Glu Leu Ile Gly Ser

515

520

525

5

Ser Gln Arg Tyr Gly His Leu Phe Glu Lys Glu Gln Ile Trp Ser Leu

530

535

540

10

Glu Pro His Ile Gln Met Tyr Leu Glu Asn Ile Ser Lys Asn Ile Leu

545

550

555

560

15 Pro Arg Asn

<210> 18

20

<211> 666

<212> DNA

25

<213> Crenarchaeote

<400> 18

30

ttgttgatc ctatggaatt taaatctaca ttggccgttt ttgatatgga tgggacgcta 60

attgatggaa ggctaattga ggtattgtca aaaaagtttg gcttgatgc tcaggtcaga 120

cacatccagt ccgacaaatc cattccaggc tatgttaaga cacagaagat agccgctgtg 180

35

attaggggaa tagaagaaag ggaaatagaa attgcttttg actccatccc ccctgcaaag 240

aacagccagg aggtgatatc tttgctgaag aaaaaagggt tcagaatagg gataattaca 300

40

gatagttaca gtgttgctgc tcaggccttg gtgaacaaac ttgatttgga ctttttttat 360

gcaaataaat tgaaggtaga caatgggata gtcaccggag aaataaatat gccgttagga 420

tgggaaaaaa tagactgttt ttgcaagaat tctgtgtgta agagatatca catggaaatc 480

catgcaaaga aaatctgtgc agacataaaa-aatacaattg ctattggcga tactaaaggt 540

5

gacctgtgca tgataaagca ggcaggaata ggtatcgcat atatgcctaa ggataaatat 600

ataaatgaaa caataaataa ggtaaacaca ccggatatga ttggtgtcct tgattttata 660

10

gagtag 666

<210> 19

15

<211> 221

<212> PRT

<213> Crenarchaeote

20

<400> 19

25

Met Leu Tyr Pro Met Glu Phe Lys Ser Thr Leu Ala Val Phe Asp Met
1 5 10 15

30

Asp Gly Thr Leu Ile Asp Gly Arg Leu Ile Glu Val Leu Ser Lys Lys
20 25 30

35

Phe Gly Leu Tyr Ala Gln Val Arg His Ile Gln Ser Asp Lys Ser Ile
35 40 45

Pro Gly Tyr Val Lys Thr Gln Lys Ile Ala Ala Val Ile Arg Gly Ile
50 55 60

40

Glu Glu Arg Glu Ile Glu Ile Ala Leu Asp Ser Ile Pro Pro Ala Lys
65 70 75 80

Asn Ser Gln Glu Val Ile Ser Leu Leu Lys Lys Lys Gly Phe Arg Ile
 85 90 95

5

Gly Ile Ile Thr Asp Ser Tyr Ser Val Ala Ala Gln Ala Leu Val Asn
 100 105 110

10

Lys Leu Asp Leu Asp Phe Phe Tyr Ala Asn Glu Leu Lys Val Asp Asn
 115 120 125

15

Gly Ile Val Thr Gly Glu Ile Asn Met Pro Leu Gly Trp Glu Lys Ile
 130 135 140

20

Asp Cys Phe Cys Lys Asn Ser Val Cys Lys Arg Tyr His Met Glu Ile
 145 150 155 160

25

His Ala Lys Lys Ile Cys Ala Asp Ile Lys Asn Thr Ile Ala Ile Gly
 165 170 175

30

Asp Thr Lys Gly Asp Leu Cys Met Ile Lys Gln Ala Gly Ile Gly Ile
 180 185 190

Ala Tyr Met Pro Lys Asp Lys Tyr Ile Asn Glu Thr Ile Asn Lys Val
 195 200 205

35

Asn Thr Pro Asp Met Ile Gly Val Leu Asp Phe Ile Glu
 210 215 220

<210> 20

40

<211> 1212

<212> DNA

<213> Crenarchaeote

5

<400> 20

	atgagattag attatccacc taactatacc gagaggatag gagcagttag tatccatgcg	60
10	cttcaaaaga tttatgagat cgattccgga aagatgccca agtttaatgg cctgcatcag	120
	catcagtcta taaaggcctt tggttatgac gaactgtcaa gcatattcca agaacttgcc	180
15	atagtcattc cagtaaagaa cgaaaaaatc agccttcttg aaggagtatt gagcgggtatt	240
	ccaaatgaat gtctcatcat catagtttcc aatagccaaa ggactcctgt cgacagattt	300
	gccatggagg ttgaaatggg aaggcagtac tctagttttg cagacaagaa aataatgatt	360
20	attcaccaaa atgatcctga gctggctaata acttttaaga aaataaagta tagatccatc	420
	ctcaacacca aaagtcaggt tcgtagtgga aaggctgaag gaatgataat tggaatattg	480
25	ctggcaaaaa tgcacctaaa agagtacatt ggattttattg acagtgataa ttattttcca	540
	ggagcagtaa atgaatatgt caagatcttt gcagcgggat ttggaatggc aaccacccca	600
	tacagcaata tcagaatata gtggcggttc aaacccaaaa tcgtaaacaa ctactacaa	660
30	ttccaagat ggggtagaat ttcagaatcc agtaacaaat acctgaacgc tctaatatcc	720
	cacatcacag ggtttgaaag ggagattatc acgactggaa atgcaggtga gcatgcatta	780
35	tccatgtccc ttgcagaaaa totcaactat tcaagcggat attcgggtga gccctatgag	840
	tttatcaaca ttttagaaaa gtttgagggt ctactcccat caaacaatcc tgacatcata	900
	gaaaagggtg tcgaaatatt tcaaataagag accaggaatc cacactttca tgaggaaaaa	960
40	ggaaatgatc atttggcagg catgatgcaa gaatctcttc tcgcaataaa caacagcaaa	1020
	atttgcaaca cagaactgac cagggaataa aatgaccatt tactcatgct tcaggtaaaa	1080

cacaataatg atatgaccaa actcaacttt aagaaaaaac accttataat ggatcccata 1140

.....aaaataatac ccacgcacaa attcgccgaa tttgtagtta agaattctaa aaccttcatt 1200.....

5

agaattggat aa

1212

<210> 21

10

<211> 403

<212> PRT

15

<213> Crenarchaeote

<400> 21

20

Met	Arg	Leu	Asp	Tyr	Pro	Pro	Asn	Tyr	Thr	Glu	Arg	Ile	Gly	Ala	Val
1				5					10					15	

25

Ser	Ile	His	Ala	Leu	Gln	Lys	Ile	Tyr	Glu	Ile	Asp	Ser	Gly	Lys	Met
			20					25					30		

30

Pro	Lys	Phe	Asn	Gly	Leu	His	Gln	His	Gln	Ser	Ile	Lys	Ala	Phe	Gly
		35					40					45			

35

Tyr	Asp	Glu	Leu	Ser	Ser	Ile	Phe	Gln	Glu	Leu	Ala	Ile	Val	Ile	Pro
	50					55					60				

40

Val	Lys	Asn	Glu	Lys	Ile	Ser	Leu	Leu	Glu	Gly	Val	Leu	Ser	Gly	Ile
65					70					75					80

Pro	Asn	Glu	Cys	Leu	Ile	Ile	Ile	Val	Ser	Asn	Ser	Gln	Arg	Thr	Pro
				85						90				95	

Val Asp Arg Phe Ala Met Glu Val Glu Met Val Arg Gln Tyr Ser Ser

100

105

110

5

Phe Ala Asp Lys Lys Ile Met Ile Ile His Gln Asn Asp Pro Glu Leu

115

120

125

10

Ala Asn Thr Phe Lys Lys Ile Lys Tyr Arg Ser Ile Leu Asn Thr Lys

130

135

140

15

Ser Gln Val Arg Ser Gly Lys Ala Glu Gly Met Ile Ile Gly Ile Leu

145

150

155

160

20

Leu Ala Lys Met His Leu Lys Glu Tyr Ile Gly Phe Ile Asp Ser Asp

165

170

175

25

Asn Tyr Phe Pro Gly Ala Val Asn Glu Tyr Val Lys Ile Phe Ala Ala

180

185

190

Gly Phe Gly Met Ala Thr Thr Pro Tyr Ser Asn Ile Arg Ile Ser Trp

195

200

205

30

Arg Ser Lys Pro Lys Ile Val Asn Asn Ser Leu Gln Phe Pro Arg Trp

210

215

220

35

Gly Arg Ile Ser Glu Ser Ser Asn Lys Tyr Leu Asn Ala Leu Ile Ser

225

230

235

240

40

His Ile Thr Gly Phe Glu Arg Glu Ile Ile Thr Thr Gly Asn Ala Gly

245

250

255

Glu His Ala Leu Ser Met Ser Leu Ala Glu Asn Leu Asn Tyr Ser Ser
 260 265 270

5 Gly Tyr Ser Val Glu Pro Tyr Glu Phe Ile Asn Ile Leu Glu Lys Phe
 275 280 285

10 Gly Gly Leu Leu Pro Ser Asn Asn Pro Asp Ile Ile Glu Lys Gly Ile
 290 295 300

15 Glu Ile Phe Gln Ile Glu Thr Arg Asn Pro His Phe His Glu Glu Lys
 305 310 315 320

Gly Asn Asp His Leu Ala Gly Met Met Gln Glu Ser Leu Leu Ala Ile
 325 330 335

20 Asn Asn Ser Lys Ile Cys Asn Thr Glu Leu Thr Arg Glu Ile Asn Asp
 340 345 350

25 His Leu Leu Met Leu Gln Val Lys His Asn Asn Asp Met Thr Lys Leu
 355 360 365

30 Asn Phe Lys Lys Lys His Leu Ile Met Asp Pro Ile Lys Ile Ile Pro
 370 375 380

35 Ile Asp Lys Phe Ala Glu Phe Val Val Lys Asn Ser Lys Thr Phe Ile
 385 390 395 400

Arg Ile Gly

40

<210> 22

<211> 1164

<212> DNA

5 <213> Crenarchaeote

<400> 22

10 atgagtgatg ctatcgaaaa tgtcctgatc cttcagggag gaggatcttt ggggtgcattt 60

ggttgcgggg tctacaaagc actagtaaac aataacataa aacttgatat cctgtctggc 120

15 acatcaattg gcggtttgaa tgccacagtt attgccggca gtaaagaaga tcgtccagaa 180

aaatcattgg agaatttttg gatggaaata gctgatacta ataatggtaa tattaatata 240

taccttaatt tccccttttt tgaaagtccc ttctctgggc aaattccttt ccccttgga 300

20 tcagaatcaa cactatcatt ctacagctct gccatttatg gaaatagaaa aatctttctg 360

ccaagatggg gacctgaaaa tatctttaa gatccacagt atttcacacc tagcaaatgg 420

25 acatatttgt atgaccattc acctttggta aaaaccttg aaaagtacat tgattatagc 480

aaattacagc caaacggtaa gcccaacgca aggctaataa taaccgcagt taacgtgatg 540

acggcgggagc cccttatttt tgacagtgcc aagcaacaaa taacccaaa acacatactt 600

30 gcaaccactg cctatccaac atattttttt caatgggtgg aattggaaaa agggcttttt 660

gcctgggatg gaagtttact aagcaatacc ccgctaagag aagtaataga cgcacgccc 720

35 gcaaaggaca aaagaatctt tcttgtcgag aactatcta aaaatattga aaagcttccg 780

tcaaacctac aggaagtcaa gcatagggca agagacataa tgttcagcga caagaccgtc 840

cacagtatac acatgtccaa agcaattacc cttcaactta agcttattga tgatctgtat 900

40 aaaatgctag agtattactt taattcagaa aaaatcgagg aaaaggagaa gtttgaaaaa 960

attcgtgcga gatacaaaaa agtttcagaa gaacacggcg cagagattaa aggtgtctac 1020

tatataacac gggacgagcc atccccctcc ctttatgaga atgcagactt ttcaaaaaat 1080

gcaataaagg catcgattaa tgatggagaa caaaaggctg acaggataat aaaagaaatc 1140

5

caaacgaaag gaaaacgaaa ataa 1164

<210> 23

10

<211> 387

<212> PRT

15

<213> Crenarchaeote

<400> 23

20

Met Ser Asp Ala Ile Glu Asn Val Leu Ile Leu Gln Gly Gly Gly Ser
1 5 10 15

25

Leu Gly Ala Phe Gly Cys Gly Val Tyr Lys Ala Leu Val Asn Asn Asn
20 25 30

30

Ile Lys Leu Asp Ile Leu Ser Gly Thr Ser Ile Gly Gly Leu Asn Ala
35 40 45

35

Thr Val Ile Ala Gly Ser Lys Glu Asp Arg Pro Glu Lys Ser Leu Glu
50 55 60

Asn Phe Trp Met Glu Ile Ala Asp Thr Asn Asn Gly Asn Ile Asn Thr
65 70 75 80

40

Tyr Leu Asn Phe Pro Phe Phe Glu Ser Pro Phe Pro Gly Gln Ile Pro
85 90 95

Phe Pro Leu Ala Ser Glu Ser Thr Leu Ser Phe Tyr Ser Ser Ala Ile
 100 105 110

5

Tyr Gly Asn Arg Lys Ile Phe Leu Pro Arg Trp Gly Pro Glu Asn Ile
 115 120 125

10

Phe Lys Asp Pro Gln Tyr Phe Thr Pro Ser Lys Trp Thr Tyr Leu Tyr
 130 135 140

15

Asp His Ser Pro Leu Val Lys Thr Leu Glu Lys Tyr Ile Asp Tyr Ser
 145 150 155 160

20

Lys Leu Gln Pro Asn Gly Lys Pro Asn Ala Arg Leu Ile Ile Thr Ala
 165 170 175

25

Val Asn Val Met Thr Ala Glu Pro Leu Ile Phe Asp Ser Ala Lys Gln
 180 185 190

30

Gln Ile Thr Pro Lys His Ile Leu Ala Thr Thr Ala Tyr Pro Thr Tyr
 195 200 205

Phe Phe Gln Trp Val Glu Leu Glu Lys Gly Leu Phe Ala Trp Asp Gly
 210 215 220

35

Ser Leu Leu Ser Asn Thr Pro Leu Arg Glu Val Ile Asp Ala Ser Pro
 225 230 235 240

40

Ala Lys Asp Lys Arg Ile Phe Leu Val Glu Asn Tyr Pro Lys Asn Ile
 245 250 255

Glu Lys Leu Pro Ser Asn Leu Gln Glu Val Lys His Arg Ala Arg Asp
 260 265 270

5 Ile Met Phe Ser Asp Lys Thr Val His Ser Ile His Met Ser Lys Ala
 275 280 285

10 Ile Thr Leu Gln Leu Lys Leu Ile Asp Asp Leu Tyr Lys Met Leu Glu
 290 295 300

15 Tyr Tyr Phe Asn Ser Glu Lys Ile Glu Glu Lys Glu Lys Phe Glu Lys
 305 310 315 320

Ile Arg Ala Arg Tyr Lys Lys Val Ser Glu Glu His Gly Ala Glu Ile
 325 330 335

20 Lys Gly Val Tyr Tyr Ile Thr Arg Asp Glu Pro Ser Pro Ser Leu Tyr
 340 345 350

25 Glu Asn Ala Asp Phe Ser Lys Asn Ala Ile Lys Ala Ser Ile Asn Asp
 355 360 365

30 Gly Glu Gln Lys Ala Asp Arg Ile Ile Lys Glu Ile Gln Thr Lys Gly
 370 375 380

35 Lys Arg Lys
 385

<210> 24

<211> 882

40

<212> DNA

<213> Crenarchaeote

5 <400> 24
atggaactta acgcagcagt aattgtgaaa ctcgagccgg atttttctga agggaatgta 60
agctataatt cagacggaac acttaacaga gcagaaacaa aaaacatttt ggggccccat 120
10 agcgagcag catccctagc agccctgtac tcaaaagtaa aacatggaac gcatgtttct 180
gtgggcacaa tgggtcctcc aatagcagaa tcggccttac agcaatctca actgatttgc 240
15 gacgctgatg aactgcatct ttatagtgat cgcattcttg caggagccga caccctggcc 300
acagctgaag ttttgatagc aggaataaaa aaaatggcaa atgggtcaaga tgtggacatt 360
gttttctcag ggcacagggc atctgatggc gaaacagggc aaacaggacc ccagacagca 420
20 tggaaattag gttatccgtt ccttggaat gttattgatt acgatattga cgttgtgaag 480
agaattgtaa ggttacaacg tctaatacag atttacggtc atcctgatat tatagaggag 540
atggaggcgc ctctaccggt ttttatcaca ctggacccat cctacaatcc gtcttttaac 600
25 acggtatccc aaaggctcag actagcacga aaactacagg aagcccatga tagatcacia 660
aggtataagg aatatctcaa aactttcaat gccatggaac tagaagtcaa tccaaagtct 720
30 gtccgactgc ctggctctcc caccatagtt tataaagttg aaaaaatacc aagggcaaag 780
gcaaatagaa aagcagatgt tgtggatggg tctaaccagg atagtctaag gcaggttgca 840
cgccgaatcc atgatgtttt aggggggtgta gtcataaagt ga 882
35

<210> 25

<211> 293

40 <212> PRT

<213> Crenarchaeote

5 <400> 25

Met Glu Leu Asn Ala Ala Val Ile Val Lys Leu Glu Pro Asp Phe Ser
 1 5 10 15

10

Glu Gly Asn Val Ser Tyr Asn Ser Asp Gly Thr Leu Asn Arg Ala Glu
 20 25 30

15

Thr Lys Asn Ile Leu Gly Pro His Ser Ala Ala Ala Ser Leu Ala Ala
 35 40 45

20

Leu Tyr Ser Lys Val Lys His Gly Thr His Val Ser Val Gly Thr Met
 50 55 60

25

Gly Pro Pro Ile Ala Glu Ser Ala Leu Gln Gln Ser Gln Leu Ile Cys
 65 70 75 80

30

Asp Ala Asp Glu Leu His Leu Tyr Ser Asp Arg Ile Phe Ala Gly Ala
 85 90 95

Asp Thr Leu Ala Thr Ala Glu Val Leu Ile Ala Gly Ile Lys Lys Met
 100 105 110

35

Ala Asn Gly Gln Asp Val Asp Ile Val Phe Ser Gly His Arg Ala Ser
 115 120 125

40

Asp Gly Glu Thr Gly Gln Thr Gly Pro Gln Thr Ala Trp Lys Leu Gly
 130 135 140

Tyr Pro Phe Leu Gly Asn Val Ile Asp Tyr Asp Ile Asp Val Val Lys
 145 150 155 160

5 Arg Ile Val Arg Val Gln Arg Leu Ile Lys Ile Tyr Gly His Pro Asp
 165 170 175

10 Ile Ile Glu Glu Met Glu Ala Pro Leu Pro Val Phe Ile Thr Leu Asp
 180 185 190

15 Pro Ser Tyr Asn Pro Ser Phe Asn Thr Val Ser Gln Arg Leu Arg Leu
 195 200 205

Ala Arg Asn Leu Gln Glu Ala His Asp Arg Ser Gln Arg Tyr Lys Glu
 210 215 220

20 Tyr Leu Lys Thr Phe Asn Ala Met Glu Leu Glu Val Asn Pro Lys Ser
 225 230 235 240

25 Val Gly Leu Pro Gly Ser Pro Thr Ile Val Tyr Lys Val Glu Lys Ile
 245 250 255

30 Pro Arg Ala Lys Ala Asn Arg Lys Ala Asp Val Val Asp Gly Ser Asn
 260 265 270

35 Gln Asp Ser Leu Arg Gln Val Ala Arg Arg Ile His Asp Val Leu Gly
 275 280 285

Gly Val Val Ile Lys
 290

40
 <210> 26

<211> 1284

<212> DNA

5 <213> Crenarchaeote

<400> 26

10 gtgacatcat cactatctgc catacctgac gctaaactag acgaaaggcc aaacccaaat 60
 gcccatgtta atgacaaccc agaaaaagaa aggggagaca acaacaggca tctgtatggt 120
 gtgatagaac aagaggaagg caccatatta cctgtgagtt ttgaaatgct tggtgaggca 180
 15 agaaggctaa tggatgattt taatcacaaa tacaagccag aggaaaaagt ggttgcgatt 240
 atactcggcc ataacatcaa gcacctgtgc caggaactaa tccaccatgg tgcagacgca 300
 20 gtgatttatg ccgaccaccc ggagctccgc caccaagaa atcttcttta taaaagggtt 360
 gtctgcaaaa ttgctacgga caaagagagc gccgccagaa tttggccatc aaatcccgat 420
 ttttaacagac cccgttacat gtttttttcc gcagatgaca caggaaggca tttatcatca 480
 25 accgttttgg cagaattgca atcagggctg gcatcagaca taaacaaact tgttatcaat 540
 gatttagaaa taaggcatga acacaagaca aagggtaaac ccattgtcta tgaaaagaca 600
 30 ottgaaatgt acagaccaga cttttcaggc tttctttgga ccaccatact ctgcttggat 660
 aatataaatc ccgagaacag aaggaaattc catccacagg catgcagtat aatcccaggc 720
 gtctttcccc aaatggaagg agatacggat agaaagggtta ccataataga gttcagccca 780
 35 accatagccc aggaagacct tagaataaaa ataatcaaca gaagagtaat caaaagcaaa 840
 gtcgatttta gcaataaaaa aataatcggt agttttggaa ggggaataaa ggagtctccc 900
 40 gaacaaaaca taaaactgat agagaacctt gcaaaggaaa tagaagcaga aataggaata 960
 tcactgccca tttcaaagaa accctatcca ataagcgaaa gtctgtcgtc aacctatatg 1020

attcctgaca gggttatcgg cacaagcgga agaaaggtaa atcctcaggt gtattttgca 1080

ataggaataa gcggggctgt ccaacacata gccgggatga aagaatcgga atttgtgatt 1140

5

tccatcaatc cagacagtga agctcccata atagatgaat ccgatgtttt aatcaaagga 1200

aaaatcgagc aggtgctgcc tctcctgata aatgaattaa aaaaatacaa agagagactg 1260

10

caaataccac aggagataga atga 1284

<210> 27

15

<211> 427

<212> PRT

<213> Crenarchaeote

20

<400> 27

25

Met Thr Ser Ser Leu Ser Ala Ile Pro Asp Ala Lys Leu Asp Glu Arg
1 5 10 15

30

Pro Asn Gln Asn Ala His Val Asn Asp Asn Pro Glu Lys Glu Arg Gly
20 25 30

35

Asp Asn Asn Arg His Leu Tyr Val Val Ile Glu Gln Glu Glu Gly Thr
35 40 45

40

Ile Leu Pro Val Ser Phe Glu Met Leu Gly Glu Ala Arg Arg Leu Met
50 55 60

Asp Asp Phe Asn His Lys Tyr Lys Pro Glu Glu Lys Val Val Ala Ile
65 70 75 80

Ile Leu Gly His Asn Ile Lys His Leu Cys Gln Glu Leu Ile His His

85

90

95

5

Gly Ala Asp Ala Val Ile Tyr Ala Asp His Pro Glu Leu Arg His Pro

100

105

110

10

Arg Asn Leu Leu Tyr Thr Lys Val Val Cys Gln Ile Ala Thr Asp Lys

115

120

125

15

Glu Ser Ala Ala Arg Ile Trp Pro Ser Asn Pro Asp Phe Asn Arg Pro

130

135

140

20

Arg Tyr Met Phe Phe Ser Ala Asp Asp Thr Gly Arg His Leu Ser Ser

145

150

155

160

Thr Val Leu Ala Glu Leu Gln Ser Gly Leu Ala Ser Asp Ile Asn Lys

165

170

175

25

Leu Val Ile Asn Asp Leu Glu Ile Arg His Glu His Lys Thr Lys Gly

180

185

190

30

Lys Pro Ile Val Tyr Glu Lys Thr Leu Glu Met Tyr Arg Pro Asp Phe

195

200

205

35

Ser Gly Phe Leu Trp Thr Thr Ile Leu Cys Leu Asp Asn Ile Asn Pro

210

215

220

40

Glu Asn Arg Arg Lys Phe His Pro Gln Ala Cys Ser Ile Ile Pro Gly

225

230

235

240

Val Phe Pro Gln Met Glu Gly Asp Thr Asp Arg Lys Gly Thr Ile Ile
 245 250 255

5 Glu Phe Ser Pro Thr Ile Ala Gln Glu Asp Leu Arg Ile Lys Ile Ile
 260 265 270

10 Asn Arg Arg Val Ile Lys Ser Lys Val Asp Phe Ser Asn Lys Lys Ile
 275 280 285

15 Ile Val Ser Phe Gly Arg Gly Ile Lys Glu Ser Pro Glu Gln Asn Ile
 290 295 300

Lys Leu Ile Glu Asn Leu Ala Lys Glu Ile Glu Ala Glu Ile Gly Ile
 305 310 315 320

20 Ser Leu Pro Ile Ser Lys Lys Pro Tyr Pro Ile Ser Glu Ser Leu Ser
 325 330 335

25 Ser Thr Tyr Met Ile Pro Asp Arg Val Ile Gly Thr Ser Gly Arg Lys
 340 345 350

30 Val Asn Pro Gln Val Tyr Phe Ala Ile Gly Ile Ser Gly Ala Val Gln
 355 360 365

35 His Ile Ala Gly Met Lys Glu Ser Glu Phe Val Ile Ser Ile Asn Pro
 370 375 380

Asp Ser Glu Ala Pro Ile Ile Asp Glu Ser Asp Val Leu Ile Lys Gly
 385 390 395 400

40 Lys Ile Glu Gln Val Leu Pro Leu Leu Ile Asn Glu Leu Lys Lys Tyr
 405 410 415

Lys Glu Arg Leu Gln Ile Pro Gln Glu Ile Glu

420

425

5

<210> 28

<211> 1878

10

<212> DNA

<213> Crenarchaeote

15

<400> 28

atgacaatgg aaagttttga tgtggcgata attgggtggag ggtctgctgg acttgcgcca 60

20

cttgagcacc totccaattt gggaaaacag gcaatcctca tagaggcagg aaaaaaata 120

ggaacaaaa acgtgtctgg gggcatattg tattccaaaa aaacagcaac tggaaaggctc 180

cacaatgtag aagatgtgtt tgataatttt ctggcagacg ctccgctgga aaggaagata 240

25

ataaaataca tgcttcacgc cgtctcaagg gaaaaagcgt tctctctgga cctgactttg 300

gcacacgact atcaaacgaa ttttgggtac accgtcctgc tcaacaaact actttcatgg 360

30

tttgcaaggg aagcatctca aagtgcagaa aaactgggtg gagggataat aacagggtgtc 420

catttaaggt cgataatctg gaaagatgac agtaccataa ttatagagac agatgaactt 480

gagccgttcc aggtaaaggc agtcattgca gctgaacggg ttaactcaga ggttgcgcaa 540

35

ataacagggtg ccagaagcaa gttcacaccg tctgacctct accagggcgt aaaggtgggtg 600

gcaaaattac cagagggggtt gcttgaagag agattcgggg tctcggaaaa cgagggagcg 660

40

gtcaccttt tttcaggcga cataacgcta aaccacattg gaggagggtt cctttacaca 720

aacagggaca ccatctcaat tggcgcagta taccattatg actctctaata tgaaaagcct 780

acagagccca atgcgctggt caatgcgtta ctgtcaaata cgtttgtgat ggaattgata 840

aaggacgagg ttccaaggat caaggaggac tacagggatc ttccaāaggā tgāagaacta 900

5
aggattaggt tcaaatacaa taaattgata aaaagctgga atgaactaca ccacacatat 960

tattcaccat ctgccgttgc agagcttgtg gcgcagggaa aatacaaata aagggaggag 1020

10
atcaaggaca aaattgattc attgtacaat gagcttgtaa caaaatacaa cacagaattt 1080

gaaacaaatt acgtggagtt agagtacagc gccaaactgg ttccagatgg aaaaagggtgc 1140

15
agaatgaaaa aaccctactt taaaaacata ttatttgtcg gtgatgctgc gggcaggggc 1200

attttccttg ggccacgcac agagggcctc aacgtaggca ttgatgacgc ggttagggcc 1260

gcagaagctg tctcaaagtc aatagatcaa aataactttc agtttgacaa cattggtgaa 1320

20
cgctacacta aatcagtgga tgaaagtcca tataccgcag acatgagcag gatcgacgca 1380

aactatctca aagccgttct tgattgcaca aaaaagggtc caaaaaacac tcttgggttt 1440

aagtatgggt ctattgtcaa attgatgtca aatagcacct ttaggaatgt atccatagga 1500

25
attgcaaact ctatagggtg caaaaggctt ttacctgtga ttgagtcaga caaacctac 1560

aatcaaattc ccatcgagat tgcggagaga aatggcaaag atttgcgga aagctattcc 1620

30
atagagattc ccaccattgc cgagcgtatt gctaacttga actataatga cgattcactg 1680

tcacacatca aggttttgaa ctgcgaaagt gactttatga aaaaaatggt ccaactgtgc 1740

cctaccaaata gctacagtat tgagaatgag cggataatgc tacagcacga aggatgcata 1800

35
gagtgtggga catgcgcaag agaaacagaa tggaggcatc ctggtgggga aaaaggaata 1860

atctataatt acgggtaa 1878

40

<210> 29

<211> 625

<212> PRT

5 <213> Crenarchaeote

<400> 29

10

Met Thr Met Glu Ser Phe Asp Val Ala Ile Ile Gly Gly Gly Ser Ala
 1 5 10 15

15

Gly Leu Ala Ala Leu Glu His Leu Ser Asn Leu Gly Lys Gln Ala Ile
 20 25 30

20

Leu Ile Glu Ala Gly Lys Lys Ile Gly Thr Lys Asn Val Ser Gly Gly
 35 40 45

25

Ile Leu Tyr Ser Lys Lys Thr Ala Thr Gly Lys Val His Asn Val Glu
 50 55 60

30

Asp Val Phe Asp Asn Phe Leu Ala Asp Ala Pro Leu Glu Arg Lys Ile
 65 70 75 80

35

Ile Lys Tyr Met Leu His Ala Val Ser Arg Glu Lys Ala Phe Ser Leu
 85 90 95

40

Asp Leu Thr Leu Ala His Asp Tyr Gln Thr Asn Phe Gly Tyr Thr Val
 100 105 110

Leu Leu Asn Lys Leu Leu Ser Trp Phe Ala Arg Glu Ala Ser Gln Ser
 115 120 125

Ala Glu Lys Leu Gly Gly Gly Ile Ile Thr Gly Val His Leu Arg Ser
 130 135 140

5 Ile Ile Trp Lys Asp Asp Ser Thr Ile Ile Ile Glu Thr Asp Glu Leu
 145 150 155 160

10 Glu Pro Phe Gln Val Lys Ala Val Ile Ala Ala Asp Gly Val Asn Ser
 165 170 175

15 Glu Val Ala Gln Ile Thr Gly Ala Arg Ser Lys Phe Thr Pro Ser Asp
 180 185 190

Leu Tyr Gln Gly Val Lys Val Val Ala Lys Leu Pro Glu Gly Leu Leu
 195 200 205

20 Glu Glu Arg Phe Gly Val Ser Glu Asn Glu Gly Ala Ala His Leu Phe
 210 215 220

25 Ser Gly Asp Ile Thr Leu Asn His Ile Gly Gly Gly Phe Leu Tyr Thr
 225 230 235 240

30 Asn Arg Asp Thr Ile Ser Ile Gly Ala Val Tyr His Tyr Asp Ser Leu
 245 250 255

35 Ile Glu Lys Pro Thr Glu Pro Asn Ala Leu Val Asn Ala Leu Leu Ser
 260 265 270

Asn Pro Phe Val Met Glu Leu Ile Lys Asp Glu Val Pro Arg Ile Lys
 275 280 285

40 Glu Asp Tyr Arg Asp Leu Ser Lys Asp Glu Glu Leu Arg Ile Arg Phe
 290 295 300

Lys Ser Asn Lys Leu Ile Lys Ser Trp Asn Asp Leu His His Thr Tyr
 305 310 315 320

5

Tyr Ser Pro Ser Ala Val Ala Glu Leu Val Ala Gln Gly Lys Tyr Lys
 325 330 335

10

Ser Arg Glu Glu Ile Lys Asp Lys Ile Asp Ser Leu Tyr Asn Glu Leu
 340 345 350

15

Val Thr Lys Tyr Asn Thr Glu Phe Glu Thr Asn Tyr Val Glu Leu Glu
 355 360 365

20

Tyr Ser Ala Lys Leu Val Pro Asp Gly Lys Arg Cys Arg Met Lys Lys
 370 375 380

25

Pro Tyr Phe Lys Asn Ile Leu Phe Val Gly Asp Ala Ala Gly Arg Gly
 385 390 395 400

30

Ile Phe Leu Gly Pro Arg Ile Glu Gly Leu Asn Val Gly Ile Asp Asp
 405 410 415

Ala Val Arg Ala Ala Glu Ala Val Ser Lys Ser Ile Asp Gln Asn Asn
 420 425 430

35

Phe Gln Phe Asp Asn Ile Gly Glu Arg Tyr Thr Lys Ser Val Asp Glu
 435 440 445

40

Ser Pro Tyr Thr Ala Asp Met Ser Arg Ile Asp Ala Asn Tyr Leu Lys
 450 455 460

Ala Val Leu Asp Cys Thr Lys Lys Val Pro Lys Asn Thr Leu Gly Phe
 465 470 475 480

5 Lys Tyr Gly Ser Ile Val Lys Leu Met Ser Asn Ser Thr Phe Arg Asn
 485 490 495

10 Val Ser Ile Gly Ile Ala Asn Ser Ile Gly Tyr Lys Arg Leu Leu Pro
 500 505 510

15 Val Ile Glu Ser Asp Lys Thr Tyr Asn Gln Ile Pro Ile Glu Ile Ala
 515 520 525

Glu Arg Asn Gly Lys Asp Leu Arg Lys Ser Tyr Ser Ile Glu Ile Pro
 530 535 540

20 Thr Ile Ala Glu Arg Ile Ala Asn Leu Asn Tyr Asn Asp Asp Ser Leu
 545 550 555 560

25 Ser His Ile Lys Val Leu Asn Ser Gln Ser Asp Phe Met Lys Lys Met
 565 570 575

30 Val Gln Leu Cys Pro Thr Lys Cys Tyr Ser Ile Glu Asn Glu Arg Ile
 580 585 590

35 Met Leu Gln His Glu Gly Cys Ile Glu Cys Gly Thr Cys Ala Arg Glu
 595 600 605

Thr Glu Trp Arg His Pro Arg Gly Glu Lys Gly Ile Ile Tyr Asn Tyr
 610 615 620

40 Gly
 625

<210> 30

5 <211> 2238

<212> DNA

<213> Crenarchaeote

10

<400> 30

15

ttggaagggtt cttctctaata acataataat aatctccatg acattggcac aaataatgat 60

aatgtatgtg aaaataaata cattgacgca aaccccggtt tgcgcacaa acgcgggtttt 120

ggtgacgtca acaataacga gaacgataat gatggtggtg ttgacgccgg tgcacccacc 180

20

acaaagggtgt actatggccc tgaaaacgca aacaacgccca ttttgagggtt catagacagg 240

gccaatgtga agatagactc ttgcataaac tccgtggccc cgtccgtgat gataggggtt 300

gacgccataa gggagaaaag ggttgacgcg gtcaaaaaca ggggccttaa actgcggtat 360

25

gtaaccgaaa taacaaagga caacgtcggc tatgtcaagg agatgctctc gttttcggag 420

attaggcacc tggatgggct gaagggaac tttgaggtgg ccgaccagag ggagtatgtg 480

30

gctgtcgcca cccttcacgc ggcacagtca ataccccagc ttttgttcag caacctccct 540

gagattgcag agcagcagca gtttgtgttt gacagctttt ggggcagggc gttgcccga 600

gagcacagga taaaggagct ggaggatggg gttgtcatgc ccgtctcact ggtcttctcc 660

35

aactacaagg acgcggtcca aaggaggttt gaaatgataa gcagggccaa aaggagata 720

ctgataatgt actccacggt taacgcgttc cacctgcagg aaaaaggcgg cacactgcaa 780

40

ctcttgaagg agatggtgga gcaaaacgac agcctgagga tcaacatcct cacgccgatg 840

gatgcctcag tgcgagagtc cttgtccttg aggctcctca caaaatacag gcccaacatc 900

caagtccagg acattgcgcc aagcattggc atcaagataa agacactggt tgtggacaga 960

~~aaggagtcgc tggatgatga gctgatacac gcaagggagg aggtggcaac cgccgcaatc~~ 1020

5

ggcttttcaa tctactccaa cagcgagcca acggtattgt cttactcgtc catatttgag 1080

gtcctctatg accagagtgt cctgttcag cagcttgacc aaaacgacaa agtcaaaagc 1140

10

gagttcataa acgtggctgc gcatgagttg cgaacgccaa tcatgcccat cctaaacggt 1200

gtggagatac tggaggagaa gcttggcgaa agaaaaacag agtttcagcg ggagcttgac 1260

15

atgataacaa gaaacgcgtc ccggctgcag aaccttgccg aaagcattct gcaggtgagc 1320

agaatcgaaa gcggaagctt tagcctggac atccaaaaaa atgtggatat ccacaacctg 1380

atttcccagg tgatagagga cattgagaaa aaatacgcct acaaggagaa ggcaaacaag 1440

20

gtggcgatag tgtttttgcc atctgacggc aacagaaatg gcgggtactc cagaggtggc 1500

ggcggcgcaa aggcagaagg ggtaaaggca gcggcgaggag caaaacaggc gcaaaaagag 1560

acgcagcaaa aagagcagtg ggtagaacct gtaaattggc ccaaccacct tttgtatgta 1620

25

gactgtgatc cgcaaaagat aagccagggt gttttcaacc tgctggacaa cgcaatgaag 1680

ttcaccaatg acggcaagat tgttgtttcc acggcagtgta tgggtgagtc ttctcccttc 1740

30

acttccacct ctcaggaaag tgataacctca aacactgcta cagctggtaa aggcaatggg 1800

ggcagagtgg atagcagcag cgacagcgac aacggtggtg gtgacaatgg tggtgaccac 1860

35

atcgggaggc agaaagaagg cgcggtgcta gtcacagtgc aggacaccgg ggttgggctc 1920

aactccaaaa taagggatca gctgtttcag aaatttgta caaagtcaaa ccagggaacc 1980

ggccttggcc tatacctgtc aaggaaaatt gttgaggagc atggtggaaa aatatggttt 2040

40

gaggagacaa acagcaaggc cggcaacagc agcagcagga acaacactaa agataaagac 2100

gaaggcattg atgaaatact gcaccacctt ggcagtgaag gaaaaatagg cgccacattc 2160

aaatttgatca tacctgtctc cctgccttcc catatgccga caaaagacat gccagaaaaa 2220

aacgatgaag gaaaatga 2238

5

<210> 31

<211> 745

10

<212> PRT

<213> Crenarchaeote

15

<400> 31

20

Met Glu Gly Ser Ser Leu Ile His Asn Asn Asn Leu His Asp Ile Gly
1 5 10 15

Thr Asn Asn Asp Asn Val Cys Glu Asn Lys Tyr Ile Asp Ala Asn Pro
20 25 30

25

Gly Leu Ser His Lys Arg Gly Phe Gly Asp Val Asn Asn Asn Glu Asn
35 40 45

30

Asp Asn Asp Gly Gly Val Asp Ala Gly Ala Pro Thr Thr Lys Val Tyr
50 55 60

35

Tyr Gly Pro Glu Asn Ala Asn Asn Ala Ile Leu Arg Phe Ile Asp Arg
65 70 75 80

40

Ala Asn Val Lys Ile Asp Ser Cys Ile Asn Ser Val Ala Pro Ser Val
85 90 95

Met Ile Gly Val Asp Ala Ile Arg Glu Lys Arg Val Asp Ala Val Lys
 100 105 110

5 Asn Arg Gly Leu Lys Leu Arg Tyr Val Thr Glu Ile Thr Lys Asp Asn
 115 120 125

10 Val Gly Tyr Val Lys Glu Met Leu Ser Phe Ser Glu Ile Arg His Leu
 130 135 140

15 Asp Gly Leu Lys Gly Asn Phe Glu Val Ala Asp Gln Arg Glu Tyr Val
 145 150 155 160

Ala Val Ala Thr Leu His Ala Ala Gln Ser Ile Pro Gln Leu Leu Phe
 165 170 175

20 Ser Asn Leu Pro Glu Ile Ala Glu Gln Gln Gln Phe Val Phe Asp Ser
 180 185 190

25 Phe Trp Gly Arg Ala Leu Pro Ala Glu His Arg Ile Lys Glu Leu Glu
 195 200 205

30 Asp Gly Val Val Met Pro Val Ser Leu Val Phe Ser Asn Tyr Lys Asp
 210 215 220

35 Ala Val Gln Arg Glu Phe Glu Met Ile Ser Arg Ala Lys Arg Glu Ile
 225 230 235 240

Leu Ile Met Tyr Ser Thr Val Asn Ala Phe His Leu Gln Glu Lys Gly
 245 250 255

40 Gly Thr Leu Gln Leu Leu Lys Glu Met Val Glu Gln Asn Asp Ser Leu
 260 265 270

	Arg	Ile	Asn	Ile	Leu	Thr	Pro	Met	Asp	Ala	Ser	Val	Arg	Glu	Ser	Leu		
			275					280						285				
5																		
	Ser	Leu	Arg	Leu	Leu	Thr	Lys	Tyr	Arg	Pro	Asn	Ile	Gln	Val	Gln	Asp		
		290					295					300						
10																		
	Ile	Ala	Pro	Ser	Ile	Gly	Ile	Lys	Ile	Lys	Thr	Leu	Val	Val	Asp	Arg		
	305					310					315					320		
15																		
	Lys	Glu	Ser	Leu	Val	Met	Glu	Leu	Ile	His	Ala	Arg	Glu	Glu	Val	Ala		
					325					330					335			
20																		
	Thr	Ala	Ala	Ile	Gly	Phe	Ser	Ile	Tyr	Ser	Asn	Ser	Glu	Pro	Thr	Val		
				340					345					350				
25																		
	Leu	Ser	Tyr	Ser	Ser	Ile	Phe	Glu	Val	Leu	Tyr	Asp	Gln	Ser	Val	Leu		
		355						360					365					
30																		
	Phe	Gln	Gln	Leu	Asp	Gln	Asn	Asp	Lys	Val	Lys	Ser	Glu	Phe	Ile	Asn		
		370					375					380						
35																		
	Val	Ala	Ala	His	Glu	Leu	Arg	Thr	Pro	Ile	Met	Pro	Ile	Leu	Asn	Gly		
	385					390					395					400		
40																		
	Val	Glu	Ile	Leu	Glu	Glu	Lys	Leu	Gly	Glu	Arg	Lys	Thr	Glu	Phe	Gln		
					405					410					415			
45																		
	Arg	Glu	Leu	Asp	Met	Ile	Thr	Arg	Asn	Ala	Ser	Arg	Leu	Gln	Asn	Leu		
			420						425					430				

Ala Glu Ser Ile Leu Gln Val Ser Arg Ile Glu Ser Gly Ser Phe Ser
 435 440 445

5 Leu Asp Ile Gln Lys Asn Val Asp Ile His Asn Leu Ile Ser Gln Val
 450 455 460

10 Ile Glu Asp Ile Glu Lys Lys Tyr Ala Tyr Lys Glu Lys Ala Asn Lys
 465 470 475 480

15 Val Ala Ile Val Phe Leu Pro Ser Asp Gly Asn Arg Asn Gly Gly Tyr
 485 490 495

Ser Arg Gly Gly Gly Gly Ala Lys Ala Glu Gly Val Lys Ala Ala Ala
 500 505 510

20 Gly Ala Lys Gln Ala Gln Lys Glu Thr Gln Gln Lys Glu Gln Trp Val
 515 520 525

25 Glu Pro Val Asn Gly Pro Asn His Leu Leu Tyr Val Asp Cys Asp Pro
 530 535 540

30 Gln Lys Ile Ser Gln Val Val Phe Asn Leu Leu Asp Asn Ala Met Lys
 545 550 555 560

35 Phe Thr Asn Asp Gly Lys Ile Val Val Ser Thr Ala Val Met Gly Glu
 565 570 575

Ser Ser Pro Phe Thr Ser Thr Ser Gln Glu Ser Asp Thr Ser Asn Thr
 580 585 590

40 Ala Thr Ala Gly Lys Gly Asn Gly Gly Arg Val Asp Ser Ser Ser Asp
 595 600 605

Ser Asp Asn Gly Gly Gly Asp Asn Gly Gly Asp His Ile Gly Arg Gln
 610 615 620

5

Lys Glu Gly Ala Val Leu Val Thr Val Gln Asp Thr Gly Val Gly Leu
 625 630 635 640

10

Asn Ser Lys Ile Arg Asp Gln Leu Phe Gln Lys Phe Val Thr Lys Ser
 645 650 655

15

Asn Gln Gly Thr Gly Leu Gly Leu Tyr Leu Ser Arg Lys Ile Val Glu
 660 665 670

20

Glu His Gly Gly Lys Ile Trp Phe Glu Glu Thr Asn Ser Lys Gly Gly
 675 680 685

25

Asn Ser Ser Ser Arg Asn Asn Thr Lys Asp Lys Asp Glu Gly Ile Asp
 690 695 700

30

Glu Ile Leu His His Leu Gly Ser Glu Gly Lys Ile Gly Ala Thr Phe
 705 710 715 720

Lys Phe Val Ile Pro Val Ser Leu Pro Ser His Met Pro Thr Lys Asp
 725 730 735

35

Met Pro Glu Lys Asn Asp Glu Gly Lys
 740 745

40

<210> 32

<211> 519

<212> DNA

<213> Crenarchaeote

5

<400> 32

ttgcaaagca gtcatttttc taaaataatc acaatttgca gaatgccgtc acttcatctt 60

10 gttgcatatg gtttaatttt ggatattttc gaaagcccaa tcacaagggt aaacggtaga 120

acaagtcact tgattattaa aatatatcca catatggata acaatacaag gatgagttct 180

15 ttagcaatcg agtttttttt atcccttttt tcaataacgt tactttctaa aagaatatac 240

caaccagtga aatcaaagtc atatacctac catgacaagc atccatttca gtacaagatg 300

gaggattatg caaaccacaa caaaattgta gactataaaa actgcttact tttttttcaa 360

20 gtatcgatgt tacaacacaa taaaataatt aggattcggg ttccagggtt gttttataca 420

ggtggctgga tttccctcac actaaagttt ttgatatcca catcatttgc accatcccac 480

ctgaaagtag caatgggggc tccccaggat ataattctga 519

25

<210> 33

<211> 172

30

<212> PRT

<213> Crenarchaeote

35

<400> 33

Met Gln Ser Ser His Leu Ser Lys Ile Ile Thr Ile Cys Arg Met Pro

40 1 5 10 15

Ser Leu His Leu Val Ala Tyr Gly Leu Ile Leu Asp Ile Phe Glu Ser
 20 25 30

5 Pro Ile Thr Arg Leu Asn Gly Arg Thr Ser His Leu Ile Ile Lys Ile
 35 40 45

10 Tyr Pro His Met Asp Asn Asn Thr Arg Met Ser Ser Leu Ala Ile Glu
 50 55 60

15 Phe Phe Leu Ser Leu Phe Ser Ile Thr Leu Leu Ser Lys Arg Ile Tyr
 65 70 75 80

Gln Pro Val Lys Ser Lys Ser Tyr Thr Tyr His Asp Lys His Pro Phe
 85 90 95

20 Gln Tyr Lys Met Glu Asp Tyr Ala Asn His Asn Lys Ile Val Asp Tyr
 100 105 110

25 Lys Asn Cys Leu Leu Phe Phe Gln Val Ser Met Leu Gln Lys Asn Lys
 115 120 125

30 Ile Ile Arg Ile Arg Val Pro Gly Leu Phe Tyr Thr Gly Gly Trp Ile
 130 135 140

35 Ser Leu Thr Leu Lys Phe Leu Ile Ser Thr Ser Phe Ala Pro Ser His
 145 150 155 160

Leu Lys Val Ala Met Gly Pro Pro Gln Asp Ile Ile
 165 170

40

<210> 34

<211> 1008

<212> DNA

5 <213> Crenarchaeote

<400> 34

10 acactgctaa caagcggagg cgcacctgtg cccgcaaccg taagcatgaa ttcaccaacc 60

aacacagttg cgaccctaaa cccctctgca gatttaacac ctggtgccac atacactgcc 120

15 agaataacaa ccggtgctac ggatgcgact ggtgtcccat tggctgccga caaggtatgg 180

acgttctctg ttgccccttc tggaggcgga gggacattag accagtttgg gataaccag 240

atttaccocg ataaggcggg tgggtggagaa aaatggttta tgaatatgca aaaccgaac 300

20 aatgatccac gaacaaacco acctgacatg gacctaaacc cagatggcag ttggaacgtt 360

aatgacgac aggtcagata taacgtgttt acatcatcag ggtaccatcc agaggatatt 420

25 gagacttacg atcactcggg actcgcaaca caaggatata tgcagtatcc aaatgattgg 480

aagaatgtgg aaatgacggg tattgtaagg gttaatagtg gagatgattc tgaaaatttc 540

gcttggtatg acaggggtgg taggcactat gatggcgaag gatgcgaggg ctcagcatat 600

30 aaagcagatc tattctatga tggaagggtt aggcttgcaa aagagcagtg gcatgtctcc 660

tatgtgtttt caagcactac cacaccctca ccttcggcgt ctagttttga tagattcatt 720

ggatttaaag ccatgatcta taaccaccaa ttggctggtg gtgagaccgt agtcaccact 780

35 gaaatatggg tagacagaaa cccggatagc ccgactctga agaacaattg gcaaaaggta 840

tacacattca ctgactcagg tgggttttga aatgatggtg aagagtgtgg tggtagccg 900

40 gatcagatta tctcctgggg aggccccatt gctactttca ggtgggatgg tgcaaagat 960

gtggatatca aaaacttttag tgtgaggga atccagccac ctgtataa 1008

<210> 35

5 <211> 335

<212> PRT

<213> Crenarchaeote

10

<400> 35

15 Thr Leu Leu Thr Ser Gly Gly Ala Pro Val Pro Ala Thr Val Ser Met
1 5 10 15

20 Asn Ser Pro Thr Asn Thr Val Ala Thr Leu Asn Pro Ser Ala Asp Leu
20 25 30

25 Thr Pro Gly Ala Thr Tyr Thr Ala Arg Ile Thr Thr Gly Ala Thr Asp
35 40 45

30 Ala Thr Gly Val Pro Leu Ala Ala Asp Lys Val Trp Thr Phe Ser Val
50 55 60

Ala Pro Ser Gly Gly Gly Gly Thr Leu Asp Gln Phe Gly Ile Thr Gln
65 70 75 80

35 Ile Tyr Pro Asp Lys Ala Gly Gly Gly Glu Lys Trp Phe Met Asn Met
85 90 95

40 Gln Asn Pro Asn Asn Asp Pro Arg Thr Asn Pro Pro Asp Met Asp Leu
100 105 110

Asn Pro Asp Gly Ser Trp Asn Val Asn Asp Asp Gln Val Arg Tyr Asn
 115 120 125

5 Val Phe Thr Ser Ser Gly Tyr His Pro Glu Asp Ile Glu Thr Tyr Asp
 130 135 140

10 His Ser Val Leu Ala Thr Gln Gly Tyr Met Gln Tyr Pro Asn Asp Trp
 145 150 155 160

15 Lys Asn Val Glu Met Thr Gly Ile Val Arg Val Asn Ser Gly Asp Asp
 165 170 175

Ser Glu Asn Phe Ala Trp Tyr Asp Arg Gly Gly Arg His Tyr Asp Gly
 180 185 190

20 Glu Gly Cys Glu Gly Ser Ala Tyr Lys Ala Asp Leu Phe Tyr Asp Gly
 195 200 205

25 Arg Val Arg Leu Ala Lys Glu Gln Trp His Val Ser Tyr Val Phe Ser
 210 215 220

30 Ser Thr Thr Thr Pro Ser Pro Ser Ala Ser Ser Phe Asp Arg Phe Ile
 225 230 235 240

35 Gly Phe Lys Ala Met Ile Tyr Asn His Gln Leu Ala Gly Gly Glu Thr
 245 250 255

Val Val Thr Thr Glu Ile Trp Val Asp Arg Asn Pro Asp Ser Pro Thr
 260 265 270

40 Leu Lys Asn Asn Trp Gln Lys Val Tyr Thr Phe Thr Asp Ser Gly Gly
 275 280 285

Phe Gly Asn Asp Gly Glu Glu Cys Gly Gly Glu Pro Asp Gln Ile Ile

290

295

300

5

Ser Trp Gly Gly Pro Ile Ala Thr Phe Arg Trp Asp Gly Ala Asn Asp

305

310

315

320

10

Val Asp Ile Lys Asn Phe Ser Val Arg Glu Ile Gln Pro Pro Val

325

330

335

15



Claims

- 5 1. A device for the isolation and/or purification of nucleic acid molecules comprising at least two layers, a first layer being adapted to bind and/or inactivate inhibitors of the activity of reagents or enzymes used in nucleic acid manipulation and a second layer being adapted to separate a plurality of nucleic acid molecules with respect to their size.
- 10 2. The device of claim 1, wherein said first layer is arranged above the second layer.
3. The device of claim 1 or 2, wherein said first layer is a first phase of a gel and said second layer is a second phase of said gel.
- 15 4. The device of claim 3, wherein said gel is an agarose-gel or a polyacrylamid-gel.
- 20 5. The device of any one of claims 1 to 4, wherein said first layer comprises polyvinylpyrrolidone (PVP), polyvinylpolypyrrolidone (PVPP), CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, antibodies, aptamers, lectins, nucleic acids or ion-exchanger.
- 25 6. The device of any one of claims 1 to 5, wherein said second layer is substantially free of PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides, aptamers, antibodies, lectins, nucleic acids or ion-exchanger.
- 30 7. The device of any one of claims 3 to 6, wherein the device is electrically biased to enhance flow of (a) sample(s) through the layers.
8. The device of any one of claims 3 to 7, wherein said first layer comprises

sample loading means.

9. The device of claim 8, wherein said loading means are provided in an array in
an upper portion of the first layer, defining an array of columns, each being
capable of isolating nucleic acid molecules.
10. The device of any one of claims 1, 5, 6, 7, 8 or 9, wherein said first layer is
arranged below the second layer.
11. The device of claim 10, which is a column comprising said first and said
second layer.
12. The device of claim 10 or 11, wherein said second layer is a first phase of a
column and said first layer is a second phase of said column.
13. The device of any one of claims 10 to 12, said first layer is a matrix comprising
PVP, PVPP, CTAB, EDTA, EGTA, cyclodextrins, proteins, (poly)peptides,
aptamers, antibodies, lectins, nucleic acids or ion-exchanger.
14. The device of any one of claims 10 to 13, wherein said second layer is a
matrix which is substantially free of PVPP PVP, PVPP, CTAB, EDTA, EGTA,
cyclodextrins, proteins, (poly)peptides, aptamers, antibodies, lectins, nucleic
acids or ion-exchanger.
15. The device of claims 13 or 14, wherein said matrix of said first and/or second
layer is selected from the group consisting of agarose, sepharoseTM,
sephadexTM, sephacrylTM, BioGelTM, superoseTM and acrylamid.
16. The device of any one of claims 1 to 15, wherein said nucleic acid molecule is
DNA or RNA.
17. The device of claim 16, wherein said DNA is genomic DNA.

18. The device of claim 16 or 17, wherein said nucleic acid molecule is derived from (micro)organisms of soil, sediments, water or symbiotic/parasitic consortia.

19. The device of claim 18, wherein said (micro)organisms are (micro)organisms of aquatic plancton, microbial mats, clusters, sludge flocs, or biofilms.

20. The device of claim 18 or 19, wherein said (micro)organism are isolated as consortia of coexisting species.

21. The device of any one of claims 1 to 20, wherein said nucleic acid molecules represent a fraction of the metagenome of a given habitat.

22. A method for the isolation of a nucleic acid molecule comprising applying a sample to the device as defined in any one of claims 1 to 21.

23. The method of claim 22, wherein a fraction of the metagenome is isolated from a given habitat.

24. A method for the generation of at least one gene library, comprising the steps of

- (a) isolating and/or purifying nucleic acid molecules from a sample using a device as defined in any one of claims 1 to 21 and optionally amplifying said nucleic acid molecules;
- (b) cloning the isolated and/or purified and optionally amplified nucleic acid molecules into appropriate vectors; and
- (c) transforming suitable hosts with said suitable vectors.

25. The method of claim 24, wherein said suitable hosts are selected from the group consisting of *E. coli*, *Pseudomonas* sp., *Bacillus* sp, *Streptomyces* sp, other actinomycetes, myxobacteria, yeasts and filamentous fungi.

26. A gene library obtained by the method of claim 24 or 25.

27. A gene library generated from metagenomic nucleic acid molecules from non-planctonic (micro)organisms comprising average insert sizes of at least 50 kB, of at least 55 kB, of at least 60 kB, of at least 80 kB, of at least 90 kB or of at least 100 kB.

28. A gene library generated from metagenomic nucleic acid molecules from planctonic (micro)organisms comprising average insert sizes of at least 85 kB, at least 90 kB, at least 95 kB, at least 100 kB, at least 120 kB, at least 140 kB, at least 160 kB or at least 200 kB.

29. A nucleic acid molecule comprising a DNA as depicted in SEQ ID NO: 1 or a DNA as deposited under EMBL accession number A3496176.

30. A nucleic acid molecule representing a part of the genome of a non-thermophilic crenarchaeota, whereby said nucleic acid molecule has at least one of the following features:

(a) it contains at least one ORF which encodes a polypeptide having the amino acid sequence SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35;

(b) comprises the DNA sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;

(c) it comprises portion of at least 20 nucleotides, preferably 100 nucleotides, more preferably at least 500 nucleotides which hybridise

under stringent conditions to the complementary strand of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34;

- (d) it is degenerate as a result of the genetic code with respect to the nucleic acid molecule of (c); or
- (e) it is at least 50% identical with the nucleic acid molecule of SEQ ID NO: 2, SEQ ID NO: 20 or SEQ ID NO: 30, 45% identical with the nucleic acid molecule of SEQ ID NO: 8 or SEQ ID NO: 26, 35% identical with the nucleic acid molecule of SEQ ID NO: 16, SEQ ID NO: 22 or SEQ ID NO: 24 or 30% identical with the nucleic acid molecule of SEQ ID NO: 4, SEQ ID NO: 14, SEQ ID NO: 18 or SEQ ID NO: 28;

- 31. The nucleic acid molecule of claim 29 or 30, which is DNA or RNA.
- 32. A vector comprising the nucleotide acid molecule of any one of claims 29 to 31.
- 33. A host transfected or transformed with the vector of claim 32.
- 34. A method for producing a (poly)peptide as encoded by a nucleic acid molecule of any one of claims 29 to 31, comprising culturing the host of claim 33 under suitable conditions and isolating said polypeptide from the culture.
- 35. A (poly)peptide encoded by a nucleic acid molecule of any one of claims 29 to 31 or as obtained by the method of claim 34.
- 36. The (poly)peptide of claim 35 which is glycosylated, phosphorylated, amidated and/or myristylated.
- 37. An antibody or an aptamer specifically recognizing the (poly)peptide of claim

35 or 36 or a fragment or epitope thereof.

38. The antibody of claim 37, which is a monoclonal antibody.

- 5 39. A transgenic non-human mammal whose somatic and germ cells comprise at least one gene encoding a functional polypeptide selected from the group consisting of:
- 10 (a) the polypeptide of claim 35 or 36;
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution.

EPO - Munich
69
20. Aug. 2002

Abstract

5 The present invention relates to a device for the isolation and/or purification of nucleic acid molecules suitable to bind and/or inactivate inhibitors of the activity of reagents or enzymes used for DNA manipulation and to separate a plurality of nucleic acid molecules with respect to their size. Moreover, the invention relates to a method for the isolation of a nucleic acid molecule comprising applying a sample to the device of the invention wherein said nucleic acid molecule preferably represents a fraction of the metagenome of a given habitat. Furthermore, the invention relates to a method for the generation of at least one gene library comprising nucleic acid molecules isolated by the method of the invention and to a nucleic acid molecule. 10 isolated by the method of the invention.






Figure 1A




Figure 1B




Figure 2




Figure 3

2-Phase Column-purification of DNA




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☒ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.